

1.
$$\frac{9x^2}{(x-1)^2(2x+1)} = \frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(2x+1)}$$

Find the values of the constants A , B and C .

(4)

$$9x^2 = A(x-1)(2x+1) + B(2x+1) + C(x-1)^2$$

Let $x = 1$

$$9 = \frac{3}{0} B$$

$$B = \frac{9}{0} \cdot 3$$

Let $x = -\frac{1}{2}$

$$\frac{9}{4} = \frac{9}{4} C$$

$$C = 1$$

Let $x = 0$

$$0 = -A + \frac{3}{0} + 1$$

$$A = \frac{4}{0} \cdot 4$$



2. $f(x) = \frac{1}{\sqrt{(9+4x^2)}}, \quad |x| < \frac{3}{2}$

Find the first three non-zero terms of the binomial expansion of $f(x)$ in ascending powers of x . Give each coefficient as a simplified fraction.

(6)

$$(9 + 4x^2)^{-1/2}$$

$$9^{-1/2} \left(1 + \frac{4}{9}x^2\right)^{-1/2}$$

$$\frac{1}{3} \left(1 + \frac{4}{9}x^2\right)^{-1/2}$$

$$\frac{1}{3} \left(1 + (-1/2) \left(\frac{4}{9}x^2\right) + \frac{(-1/2)(-3/2)}{2} \left(\frac{4}{9}x^2\right)^2\right)$$

$$\frac{1}{3} \left(1 - \frac{2}{9}x^2 + \frac{2}{27}x^4\right)$$

$$\frac{1}{3} - \frac{2}{27}x^2 + \frac{2}{81}x^4$$



3.




Figure 1

A hollow hemispherical bowl is shown in Figure 1. Water is flowing into the bowl. When the depth of the water is h m, the volume V m³ is given by

$$V = \frac{1}{12} \pi h^2 (3 - 4h), \quad 0 \leq h \leq 0.25$$

- (a) Find, in terms of π , $\frac{dV}{dh}$ when $h = 0.1$ (4)

Water flows into the bowl at a rate of $\frac{\pi}{800}$ m³s⁻¹.

- (b) Find the rate of change of h , in ms⁻¹, when $h = 0.1$ (2)

$$a) \quad V = \frac{1}{12} \pi h^2 (3 - 4h)$$

$$= \frac{1}{4} \pi h^2 - \frac{1}{3} \pi h^3$$

$$\frac{dV}{dh} = \frac{1}{2} \pi h - \pi h^2$$

$$\text{when } h = 0.1 \quad \frac{dV}{dh} = \frac{1}{20} \pi - \frac{1}{100} \pi$$

$$= \frac{1}{25} \pi$$

$$b) \quad \frac{dV}{dt} = \frac{\pi}{800}$$

$$\frac{dh}{dt} = \frac{dh}{dV} \cdot \frac{dV}{dt}$$

$$= \frac{25}{\pi} \cdot \frac{\pi}{800}$$

$$= \frac{1}{32} \text{ ms}^{-1}$$



4.




Figure 2

Figure 2 shows a sketch of the curve with equation $y = x^3 \ln(x^2 + 2)$, $x \geq 0$. The finite region R , shown shaded in Figure 2, is bounded by the curve, the x -axis and the line $x = \sqrt{2}$.

The table below shows corresponding values of x and y for $y = x^3 \ln(x^2 + 2)$.

x	0	$\frac{\sqrt{2}}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{3\sqrt{2}}{4}$	$\sqrt{2}$
y	0	0.0333	0.3240	1.3596	3.9210

- (a) Complete the table above giving the missing values of y to 4 decimal places. (2)
- (b) Use the trapezium rule, with all the values of y in the completed table, to obtain an estimate for the area of R , giving your answer to 2 decimal places. (3)
- (c) Use the substitution $u = x^2 + 2$ to show that the area of R is

$$\frac{1}{2} \int_2^4 (u - 2) \ln u \, du \quad (4)$$

- (d) Hence, or otherwise, find the exact area of R . (6)



Question 4 continued

$$b/ \frac{\sqrt{2}}{4} (0.0333 + 0.3240 + 1.3596 + \frac{3.9210}{2})$$

$$= 1.30 \text{ units}^2 \text{ 2dp}$$

$$c/ \int_0^{\sqrt{2}} x^3 \ln(x^2 + 2) dx \quad u = x^2 + 2$$

$$\int_2^4 x^3 \ln u \frac{dx}{du} du \quad \frac{du}{dx} = 2x$$

$$\frac{dx}{du} = \frac{1}{2x}$$

$$\int_2^4 x^3 \ln(u) \frac{1}{2x} dx$$

$$\int_2^4 \frac{1}{2} x^2 \ln u du$$

$$\frac{1}{2} \int_2^4 x(u-2) \ln u du$$

$$d/ \quad u = \ln u \quad \frac{dv}{dx} = u - 2$$

$$\frac{du}{dx} = \frac{1}{u} \quad v = \frac{1}{2} u^2 - 2u$$

$$= \frac{1}{2} \left[\left(\frac{1}{2} u^2 - 2u \right) \ln(u) - \int \frac{1}{u} \left(\frac{1}{2} u^2 - 2u \right) du \right]_2^4$$

$$\frac{1}{2} \left[\left(\frac{1}{2} u^2 - 2u \right) \ln u - \int \frac{1}{2} u - 2 du \right]_2^4$$

$$\frac{1}{2} \left[\left(\frac{1}{2} u^2 - 2u \right) \ln u - \left(\frac{1}{4} u^2 - 2u \right) \right]_2^4$$

$$\frac{1}{2} \left(\left[\left(\frac{1}{2} (4)^2 - 2(4) \right) \ln(4) - \left(\frac{1}{4} (4)^2 - 2(4) \right) \right] - \left[\left(\frac{1}{2} (2)^2 - 2(2) \right) \ln(2) - \left(\frac{1}{4} (2)^2 - 2(2) \right) \right] \right)$$

$$\frac{1}{2} (4 - (-2 \ln 2 + 3))$$

$$\frac{1}{2} (1 + 2 \ln 2)$$

$$\underline{\underline{\frac{1}{2} + \ln 2}}$$

Q4

(Total 15 marks)



5. Find the gradient of the curve with equation

$$\ln y = 2x \ln x, \quad x > 0, y > 0$$

at the point on the curve where $x = 2$. Give your answer as an exact value.

(7)

$$\frac{1}{y} \frac{dy}{dx} = 2 \ln x + 2$$

$$u = 2x \quad v = \ln x$$

$$\frac{du}{dx} = 2 \quad \frac{dv}{dx} = \frac{1}{x}$$

when $x = 2$

$$\ln y = 4 \ln 2$$

$$\ln y = \ln 16$$

$$y = 16$$

$$\frac{dy}{dx} = 16(2 \ln 2 + 2)$$

$$= \underline{\underline{32 \ln 2 + 32}}$$



6. With respect to a fixed origin O , the lines l_1 and l_2 are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} -5 \\ 15 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}$$

where λ and μ are scalar parameters.

(a) Show that l_1 and l_2 meet and find the position vector of their point of intersection A . (6)

(b) Find, to the nearest 0.1° , the acute angle between l_1 and l_2 . (3)

The point B has position vector $\begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix}$.

(c) Show that B lies on l_1 . (1)

(d) Find the shortest distance from B to the line l_2 , giving your answer to 3 significant figures. (4)

a/ $6 - \lambda = -5 + 2\mu$ (1)

$-3 + 2\lambda = 15 - 3\mu$

$-2 + 3\lambda = 3 + \mu$

(1) $\lambda = 11 - 2\mu$

(2) $-3 + 2(11 - 2\mu) = 15 - 3\mu$

$-3 + 22 - 4\mu = 15 - 3\mu$

$4 = \mu$

$\lambda = 11 - 8$

$= 3$

$l_1 = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + 3 \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix}$



Question 6 continued

$$L_2 = \begin{pmatrix} -5 \\ 15 \\ 3 \end{pmatrix} + 4 \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}$$

$$= \underline{\underline{\begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix}}}$$

b/ $a \cdot b = |a| |b| \cos \theta$

$$a \cdot b = (-1)(2) + (2)(-3) + (3)(1)$$

$$= -5$$

$$-5 = \sqrt{1^2 + 2^2 + 3^2} \sqrt{2^2 + 3^2 + 1^2} \cos \theta$$

$$-5 = \sqrt{14} \sqrt{14} \cos \theta$$

$$\cos \theta = \frac{-5}{14}$$

$$\theta = \underline{\underline{110.9^\circ}}$$

c/ $\begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix}$

i// $6 - \lambda = 5$
 $\lambda = 1$

ii// $-3 + 2 = -1$ ✓

iii// $-2 + 3 = 1$ ✓

d/ Shortest distance = perpendicular distance

point on $L_2 = \begin{pmatrix} -5 + 2\mu \\ 15 - 3\mu \\ 3 + \mu \end{pmatrix}$

$a \cdot b = 0$ [perp]

$$\vec{BP}_2 = \begin{pmatrix} -10 + 2\mu \\ 16 - 3\mu \\ 2 + \mu \end{pmatrix}$$

$$2(-10 + 2\mu) - 3(16 - 3\mu) + 1(2 + \mu) = 0$$



Question 6 continued

$$-20 + 4\mu - 48 + 9\mu + 2 + \mu = 0$$

$$14\mu - 66 = 0$$

$$\mu = \frac{33}{7}$$

$$BP_{\perp} = \begin{pmatrix} -4/7 \\ 13/7 \\ 47/7 \end{pmatrix}$$

$$\text{distance} = \sqrt{\left(\frac{-4}{7}\right)^2 + \left(\frac{13}{7}\right)^2 + \left(\frac{47}{7}\right)^2}$$

$$= \underline{\underline{6.99}} \quad 3 \text{ s.f.}$$

[It would have probably been easier to draw a triangle and do SOHCAHTOA!!!]

(Total 14 marks)

Q6



7.




Figure 3

Figure 3 shows part of the curve C with parametric equations

$$x = \tan \theta, \quad y = \sin \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point P lies on C and has coordinates $\left(\sqrt{3}, \frac{1}{2}\sqrt{3}\right)$.

(a) Find the value of θ at the point P . (2)

The line l is a normal to C at P . The normal cuts the x -axis at the point Q .

(b) Show that Q has coordinates $(k\sqrt{3}, 0)$, giving the value of the constant k . (6)

The finite shaded region S shown in Figure 3 is bounded by the curve C , the line $x = \sqrt{3}$ and the x -axis. This shaded region is rotated through 2π radians about the x -axis to form a solid of revolution.

(c) Find the volume of the solid of revolution, giving your answer in the form $p\pi\sqrt{3} + q\pi^2$, where p and q are constants. (7)

a/ $\sqrt{3} = \tan \theta$
 $\theta = \tan^{-1} \sqrt{3}$
 $= \frac{1}{3}\pi$

b/ $\frac{dx}{d\theta} = \sec^2 \theta$ $\frac{dy}{d\theta} = \cos \theta$



Question 7 continued

$$\frac{dy}{dx} = \frac{\cos \theta}{\sec^2 \theta}$$

$$= \cos^3 \theta$$

$$\theta = \frac{\pi}{3} \quad \frac{dy}{dx} = \frac{1}{8}$$

normal $m = -8$

$$y = -8x + c$$

$$\frac{1}{2}\sqrt{3} = -8(\sqrt{3}) + c$$

$$c = \frac{17}{2}\sqrt{3}$$

$$y = -8x + \frac{17}{2}\sqrt{3}$$

Crosses x when $y=0$

$$0 = -8x + \frac{17}{2}\sqrt{3}$$

$$8x = \frac{17}{2}\sqrt{3}$$

$$x = \frac{17}{16}\sqrt{3}$$

$$k = \frac{17}{16}$$

$$c/ \text{ volume} = \pi \int_{\text{⑥}}^{\text{③}} y^2 \frac{dx}{d\theta} d\theta$$

$$= \pi \int_0^{\frac{1}{3}\pi} \sin^2 \theta \sec^2 \theta d\theta$$

$$= \pi \int_0^{\frac{1}{3}\pi} \tan^2 \theta d\theta$$

$$\pi \int_0^{\frac{1}{3}\pi} \sec^2 \theta - 1 d\theta$$

$$\pi \left[\tan \theta - \theta + c \right]_0^{\frac{1}{3}\pi}$$



Question 7 continued

$$\pi \left[\left(\sqrt{3} - \frac{\pi}{3} \right) - 0 \right]$$

$$\underline{\underline{\frac{\sqrt{3}\pi - \pi^2}{3}}}$$

Q7

(Total 15 marks)



8. (a) Find $\int (4y+3)^{\frac{1}{2}} dy$

(2)

(b) Given that $y=1.5$ at $x=-2$, solve the differential equation

$$\frac{dy}{dx} = \frac{\sqrt{4y+3}}{x^2}$$

giving your answer in the form $y=f(x)$.

(6)

8a/ $\frac{2}{4} (4y+3)^{\frac{1}{2}}$
 $\frac{1}{2} (4y+3)^{\frac{1}{2}} + C$

b/ $\int (4y+3)^{-\frac{1}{2}} dy = \int x^{-2} dx$
 $\frac{1}{2} (4y+3)^{\frac{1}{2}} = -x^{-1} + C$

∴ $(-2, 1.5)$

$$\frac{1}{2} (4(1.5)+3)^{\frac{1}{2}} = -\frac{1}{-2} + C$$

$$\frac{3}{2} = \frac{1}{2} + C$$

$$C = 1$$

$$\frac{1}{2} (4y+3)^{\frac{1}{2}} = -x^{-1} + 1$$

$$(4y+3)^{\frac{1}{2}} = -\frac{2}{x} + 2$$

$$(4y+3) = \left(-\frac{2}{x} + 2\right)^2$$

$$4y + 3 = \left(-\frac{2}{x} + 2\right)^2 - 3$$

$$y = \frac{1}{4} \left(-\frac{2}{x} + 2\right)^2 - \frac{3}{4}$$

