

1 Find an equation for the tangent to the curve with equation $y = x^2 + \ln(4x - 1)$ at the point on the curve where $x = \frac{1}{2}$.

2 A curve has the equation $y = \sqrt{8 - e^{2x}}$.

The point P on the curve has y -coordinate 2.

a Find the x -coordinate of P .

b Show that the tangent to the curve at P has equation

$$2x + y = 2 + \ln 4.$$


3 A curve has the equation $y = 2x + 1 + \ln(4 - 2x)$, $x < 2$.

a Find and simplify expressions for $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.

b Find the coordinates of the stationary point of the curve.

c Determine the nature of this stationary point.

4



The diagram shows the curve with equation $y = \frac{3}{2x+1}$.

a Find an equation for the normal to the curve at the point $P(1, 1)$.

The normal to the curve at P intersects the curve again at the point Q .

b Find the exact coordinates of Q .

5 A quantity N is increasing such that at time t seconds,

$$N = ae^{kt}.$$

Given that at time $t = 0$, $N = 20$ and that at time $t = 8$, $N = 60$, find

a the values of the constants a and k ,

b the value of N when $t = 12$,

c the rate at which N is increasing when $t = 12$.

6

$$f(x) \equiv (5 - 2x^2)^3.$$


a Find $f'(x)$.

b Find the coordinates of the stationary points of the curve $y = f(x)$.

c Find the equation for the tangent to the curve $y = f(x)$ at the point with x -coordinate $\frac{3}{2}$, giving your answer in the form $ax + by + c = 0$, where a , b and c are integers.

- 7 A curve has the equation $y = 4x - \frac{1}{2}e^{2x}$.
- Find the coordinates of the stationary point of the curve, giving your answers in terms of natural logarithms.
 - Determine the nature of the stationary point.

8



The diagram shows the curve $y = f(x)$ where $f(x) = 3 \ln 5x - 2x$, $x > 0$.

- Find $f'(x)$.
 - Find the x -coordinate of the point on the curve at which the gradient of the normal to the curve is $-\frac{1}{4}$.
 - Find the coordinates of the maximum turning point of the curve.
 - Write down the set of values of x for which $f(x)$ is a decreasing function.
- 9 The curve C has the equation $y = \sqrt{x^2 + 3}$.
- Find an equation for the tangent to C at the point $A(-1, 2)$.
 - Find an equation for the normal to C at the point $B(1, 2)$.
 - Find the x -coordinate of the point where the tangent to C at A meets the normal to C at B .
- 10 A bucket of hot water is placed outside and allowed to cool. The surface temperature of the water, T °C, after t minutes is given by
- $$T = 20 + 60e^{-kt},$$
- where k is a positive constant.
- State the initial surface temperature of the water.
 - State, with a reason, the air temperature around the bucket.
- Given that $T = 30$ when $t = 25$,
- find the value of k ,
 - find the rate at which the surface temperature of the water is decreasing when $t = 40$.

11

$$f(x) \equiv x^2 - 7x + 4 \ln\left(\frac{x}{2}\right), \quad x > 0.$$

- Solve the equation $f'(x) = 0$, giving your answers correct to 2 decimal places.
 - Find an equation for the tangent to the curve $y = f(x)$ at the point on the curve where $x = 2$.
- 12 A curve has the equation $y = x^2 - \frac{8}{x-1}$.
- Show that the x -coordinate of any stationary point of the curve satisfies the equation $x^3 - 2x^2 + x + 4 = 0$.
 - Hence, show that the curve has exactly one stationary point and find its coordinates.
 - Determine the nature of this stationary point.