

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

Pearson Edexcel Level 3 GCE

Tuesday 20 June 2023

Afternoon

Paper
reference

9MA0/32



Mathematics Advanced **PAPER 32: Mechanics**

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

**Candidates may use any calculator allowed by Pearson regulations.
Calculators must not have the facility for symbolic algebra manipulation,
differentiation and integration, or have retrievable mathematical formulae
stored in them.**

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need*.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 50. There are 6 questions.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question*.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P72820A

©2023 Pearson Education Ltd.
N:1/1/1/1/1/1/1/1/1



P 7 2 8 2 0 A 0 1 2 0



Pearson

1. A car is initially at rest on a straight horizontal road.

The car then accelerates along the road with a constant acceleration of 3.2 ms^{-2}

Find

(a) the speed of the car after 5 s,

(1)

(b) the distance travelled by the car in the first 5 s.

(2)

a/ s
 $u = 0$
 $v = ?$
 $a = 3.2$
 $t = 5$

$$\begin{aligned} v &= u + at \\ &= 0 + 3.2(5) \\ &= 16 \end{aligned}$$

16 m/s

b/ $s = ut + \frac{1}{2}at^2$
 $= 0(5) + \frac{1}{2}(3.2)(5)^2$
 $= 40 \text{ m}$



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



P 7 2 8 2 0 A 0 3 2 0

(Total for Question 1 is 3 marks)

2.



A particle P has mass 5 kg.

The particle is pulled along a rough horizontal plane by a horizontal force of magnitude 28 N.

The only resistance to motion is a frictional force of magnitude F newtons, as shown in Figure 1.

(a) Find the magnitude of the normal reaction of the plane on P

(1)

The particle is accelerating along the plane at 1.4 m s^{-2}

(b) Find the value of F

(2)

The coefficient of friction between P and the plane is μ

(c) Find the value of μ , giving your answer to 2 significant figures.

(1)

$$a/ \quad R = 5g$$

$$b/ \quad F = ma$$

$$28 - Fr = 5(1.4)$$

$$Fr = 28 - 5(1.4)$$

$$= \underline{\underline{21 \text{ N}}}$$

$$c/ \quad Fr = \mu R$$

$$21 = \mu(5g)$$

$$\mu = \frac{21}{5g}$$

$$= \underline{\underline{0.43}}$$

Question 2 continued

(Total for Question 2 is 4 marks)



3. At time t seconds, where $t \geq 0$, a particle P has velocity $\mathbf{v} \text{ m s}^{-1}$ where

$$\mathbf{v} = (t^2 - 3t + 7)\mathbf{i} + (2t^2 - 3)\mathbf{j}$$

Find

- (a) the speed of P at time $t = 0$ (3)
- (b) the value of t when P is moving parallel to $(\mathbf{i} + \mathbf{j})$ (2)
- (c) the acceleration of P at time t seconds (2)
- (d) the value of t when the direction of the acceleration of P is perpendicular to \mathbf{i} (2)

a/ when $t=0$ $\mathbf{v} = 7\mathbf{i} - 3\mathbf{j}$

$$\text{speed} = \sqrt{7^2 + 3^2}$$

$$= 7.6 \text{ m s}^{-1}$$

b/ $\underbrace{(t^2 - 3t + 7)\mathbf{i} + (2t^2 - 3)\mathbf{j}}_{t^2 - 3t + 7 = k} = k\mathbf{i} + k\mathbf{j}$

$$t^2 - 3t + 7 = k \quad 2t^2 - 3 = k$$

$$t^2 - 3t + 7 = 2t^2 - 3$$


$$0 = t^2 + 3t - 10$$

$$0 = (t + 5)(t - 2)$$

$$\begin{matrix} t = -5 \\ \times \end{matrix} \quad \begin{matrix} t = 2 \\ \equiv \end{matrix}$$

c/ $\mathbf{a} = \frac{d\mathbf{v}}{dt}$

$$\mathbf{a} = (2t - 3)\mathbf{i} + 4t\mathbf{j}$$



d/ $2t - 3 = 0$
 $t = \frac{3}{2}$

Question 3 continued

(Total for Question 3 is 9 marks)



4. [In this question, \mathbf{i} and \mathbf{j} are horizontal unit vectors and position vectors are given relative to a fixed origin O]

A particle P is moving on a smooth horizontal plane.

The particle has constant acceleration $(2.4\mathbf{i} + \mathbf{j}) \text{ m s}^{-2}$

At time $t = 0$, P passes through the point A .

At time $t = 5 \text{ s}$, P passes through the point B .

The velocity of P as it passes through A is $(-16\mathbf{i} - 3\mathbf{j}) \text{ m s}^{-1}$

(a) Find the speed of P as it passes through B . (4)

The position vector of A is $(44\mathbf{i} - 10\mathbf{j}) \text{ m}$.

At time $t = T$ seconds, where $T > 5$, P passes through the point C .

The position vector of C is $(4\mathbf{i} + c\mathbf{j}) \text{ m}$.

(b) Find the value of T .

$$\mathbf{s} = \begin{pmatrix} 4 \\ c \end{pmatrix} - \begin{pmatrix} 44 \\ -10 \end{pmatrix}$$

(c) Find the value of c .

$$= \begin{pmatrix} -40 \\ c+10 \end{pmatrix}$$

(3)

(3)

a/ $s =$
 $u = -16\mathbf{i} - 3\mathbf{j}$
 $v = ?$
 $a = 2.4\mathbf{i} + \mathbf{j}$
 $t = 5$

$$v = \begin{pmatrix} -16 \\ -3 \end{pmatrix} + 5 \begin{pmatrix} 2.4 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} -16 \\ -3 \end{pmatrix} + \begin{pmatrix} 12 \\ 5 \end{pmatrix}$$

$$= \begin{pmatrix} -4 \\ 2 \end{pmatrix}$$

$$-4\mathbf{i} + 2\mathbf{j}$$

$$\text{speed} = \sqrt{4^2 + 2^2}$$

$$= 4.5 \text{ m/s}$$



Question 4 continued

$$s = -40i + (c+10)j$$

$$u = -16i - 3j$$

v

$$a = 2.4i + j$$

$$t = T$$

$$s = ut + \frac{1}{2}at^2$$

i/ $-40 = -16T + \frac{1}{2}(2.4)T^2$

$$0 = 1.2T^2 - 16T + 40$$

$$\underline{\underline{T = 10 \text{ s}}} \quad T = \frac{10}{X^3}$$

c/
j/

$$c+10 = -3(10) + \frac{1}{2}(1)(10)^2$$

$$c+10 = -30 + 50$$

$$c+10 = 20$$

$$\underline{\underline{c = 10}}$$



P 7 2 8 2 0 A 0 9 2 0

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




Question 4 continued

(Total for Question 4 is 10 marks)



5.

**Figure 2**

A small ball is projected with speed 28 m s^{-1} from a point O on horizontal ground.

After moving for T seconds, the ball passes through the point A .

The point A is 40 m horizontally and 20 m vertically from the point O , as shown in Figure 2.

The motion of the ball from O to A is modelled as that of a particle moving freely under gravity.

Given that the ball is projected at an angle α to the ground, use the model to

(a) show that $T = \frac{10}{7 \cos \alpha}$ (2)

(b) show that $\tan^2 \alpha - 4 \tan \alpha + 3 = 0$ (5)

(c) find the greatest possible height, in metres, of the ball above the ground as the ball moves from O to A . (3)

The model does not include air resistance.

(d) State one other limitation of the model. (1)

Question 5 continued

Horizontally

$$\begin{aligned} s &= 40 \\ u &= 28 \cos \alpha \\ v &= \\ a &= 0 \\ t &= T \end{aligned}$$

Vertically

$$\begin{aligned} s &= 20 \\ u &= 28 \sin \alpha \\ v &= \\ a &= -g \\ t &= T \end{aligned}$$

a) Horizontal

$$s = ut + \frac{1}{2} at^2$$

$$40 = 28 \cos \alpha \cdot T$$

$$\begin{aligned} T &= \frac{40}{28 \cos \alpha} \\ &= \frac{10}{7 \cos \alpha} \end{aligned}$$

$$\text{Vertically: } s = ut + \frac{1}{2} at^2$$

$$20 = 28 \sin \alpha \cdot T + \frac{1}{2} (-g) T^2$$

$$20 = 28 \sin \alpha \cdot \frac{10}{7 \cos \alpha} - \frac{1}{2} (9.8) \left(\frac{10}{7 \cos \alpha} \right)^2$$

$$20 = 40 \tan \alpha - \frac{10}{\cos^2 \alpha}$$

$$\frac{1}{\cos^2 \alpha} = \sec^2 \alpha$$

$$\begin{aligned} \cos^2 \alpha + \sin^2 \alpha &= 1 \\ 1 + \tan^2 \alpha &= \sec^2 \alpha \end{aligned}$$



Question 5 continued

$$20 = 40 \tan \alpha - 10 \sec^2 \alpha$$

$$20 = 40 \tan \alpha - 10(1 + \tan^2 \alpha)$$

$$20 = 40 \tan \alpha - 10 - 10 \tan^2 \alpha$$

$$2 = 4 \tan \alpha - 1 - \tan^2 \alpha$$

$$\tan^2 \alpha - 4 \tan \alpha + 3 = 0$$

c) $\tan \alpha = 3$ $\tan \alpha = 1$
 $\alpha = 71.6^\circ$ $= 45^\circ$

$$s = ?$$

$$u = 28 \sin \alpha \quad v^2 = u^2 + 2as$$

$$v = 0$$

$$a = -g$$

$$s = (28 \sin 71.6) \quad 0 = (28 \sin 71.6)^2 + 2(-9.8)s$$

t

$$19.6s = 705.6$$

$$s = 36 \text{ m}$$

d/ we could use a more accurate value
for g.



Question 5 continued

(Total for Question 5 is 11 marks)



6.




Figure 3

A rod AB has mass M and length $2a$.

The rod has its end A on rough horizontal ground and its end B against a smooth vertical wall.

The rod makes an angle θ with the ground, as shown in Figure 3.

The rod is at rest in limiting equilibrium. $F_r \text{ Max}$.

- (a) State the direction (left or right on Figure 3 above) of the frictional force acting on the rod at A . Give a reason for your answer.

(1)

The magnitude of the normal reaction of the wall on the rod at B is S .

In an initial model, the rod is modelled as being **uniform**.

Use this initial model to answer parts (b), (c) and (d).

- (b) By taking moments about A , show that

$$S = \frac{1}{2} Mg \cot \theta \quad (3)$$

The coefficient of friction between the rod and the ground is μ

Given that $\tan \theta = \frac{3}{4}$

- (c) find the value of μ (5)

- (d) find, in terms of M and g , the magnitude of the resultant force acting on the rod at A . (3)

In a new model, the rod is modelled as being **non-uniform**, with its centre of mass closer to B than it is to A .

A new value for S is calculated using this new model, with $\tan \theta = \frac{3}{4}$

- (e) State whether this new value for S is larger, smaller or equal to the value that S would take using the initial model. Give a reason for your answer.

(1)



Question 6 continued

a) right Friction has to oppose the motion There is a force acting to the left so to balance friction must be to the right.

b) moments about a

$$\cancel{Mg} \cos \theta = 2\cancel{a} \cdot S \sin \theta$$

$$Mg \cos \theta = 2S \sin \theta$$

$$Mg \cot \theta = 2S$$

$$\underline{\frac{1}{2} Mg \cot \theta = S}$$

c)

$$Fr = \frac{1}{2} Mg \cot \theta$$

$$\mu R_A = \frac{1}{2} Mg \left(\frac{4}{3}\right)$$

$$\mu R_A = \frac{2}{3} Mg$$

$$\mu Mg = \frac{2}{3} Mg$$

$$\underline{\mu = \frac{2}{3}}$$

d)


$$R_A = Mg$$

$$Fr = \frac{2}{3} Mg$$



P 7 2 8 2 0 A 0 1 7 2 0

Question 6 continued



$$\frac{2}{3} Mg$$

$$\left(\frac{2}{3}\right)^2 + (1)^2 = c^2$$

$$c = \sqrt{\left(\frac{2}{3}\right)^2 + 1^2}$$

$$= \frac{\sqrt{13}}{3}$$

$$\underline{\underline{\frac{\sqrt{13}}{3} Mg N}}$$

e) larger. When taking moments about A. The clockwise moment (weight) would be larger $\therefore S$ would need to be larger.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued



Question 6 continued

(Total for Question 6 is 13 marks)

TOTAL FOR MECHANICS IS 50 MARKS

