

1. (a) Find the binomial expansion of

$$\sqrt{1-8x}, \quad |x| < \frac{1}{8},$$

in ascending powers of x up to and including the term in x^3 , simplifying each term. (4)

(b) Show that, when $x = \frac{1}{100}$, the exact value of $\sqrt{1-8x}$ is $\frac{\sqrt{23}}{5}$. (2)

(c) Substitute $x = \frac{1}{100}$ into the binomial expansion in part (a) and hence obtain an approximation to $\sqrt{23}$. Give your answer to 5 decimal places. (3)

1a) $(1-8x)^{\frac{1}{2}}$

$$1 + \left(\frac{1}{2}\right)(-8x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)(-8x)^2}{2} + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)(-8x)^3}{6}$$

$$1 - 4x + -8x^2 - 32x^3$$

$$1 - 4x - 8x^2 - 32x^3$$

b/ $\sqrt{1-8\left(\frac{1}{100}\right)}$

$$\sqrt{1-\frac{8}{100}}$$

$$\sqrt{\frac{92}{100}}$$

$$\frac{\sqrt{92}}{10}$$

$$\frac{2\sqrt{23}}{10}$$

$$\frac{2\sqrt{23}}{10} = \frac{\sqrt{23}}{5}$$

c/ $\frac{\sqrt{23}}{5} = 1 - (4)\left(\frac{1}{100}\right) - 8\left(\frac{1}{100}\right)^2 - 32\left(\frac{1}{100}\right)^3$



Question 1 continued

$$\frac{\sqrt{23}}{5} = 0.959168$$

$$\sqrt{23} = \underline{\underline{4.79584}}$$

Q1

(Total 9 marks)



2.




Figure 1

Figure 1 shows a sketch of the curve with equation $y = x \ln x$, $x \geq 1$. The finite region R , shown shaded in Figure 1, is bounded by the curve, the x -axis and the line $x = 4$.

The table shows corresponding values of x and y for $y = x \ln x$.

x	1	1.5	2	2.5	3	3.5	4
y	0	0.608	1.386	2.291	3.296	4.385	5.545

- (a) Complete the table with the values of y corresponding to $x = 2$ and $x = 2.5$, giving your answers to 3 decimal places. (2)
- (b) Use the trapezium rule, with all the values of y in the completed table, to obtain an estimate for the area of R , giving your answer to 2 decimal places. (4)
- (c) (i) Use integration by parts to find $\int x \ln x \, dx$.
- (ii) Hence find the exact area of R , giving your answer in the form $\frac{1}{4}(a \ln 2 + b)$, where a and b are integers. (7)

b) $0.5 \left(\frac{0}{2} + 0.608 + 1.386 + 2.291 + 3.296 + 4.385 + \frac{5.545}{2} \right)$
 $= 7.37$ (2dp)

c) i) $u = \ln x \quad \frac{du}{dx} = \frac{1}{x}$
 $\frac{du}{dx} = \frac{1}{x} \quad v = \frac{1}{2}x^2$



Question 2 continued

$$\begin{aligned}\int x \ln x \, dx &= \frac{1}{2} x^2 \ln x - \int \frac{1}{x} \cdot \frac{1}{2} x^2 \, dx \\ &= \frac{1}{2} x^2 \ln x - \int \frac{1}{2} x \, dx \\ &= \frac{1}{2} x^2 \ln x - \frac{1}{4} x^2 + c\end{aligned}$$

$$\text{a) } \left[\frac{1}{2} x^2 \ln x - \frac{1}{4} x^2 \right]_1^4$$

$$\left[\frac{1}{2} (4)^2 \ln 4 - \frac{1}{4} (4)^2 \right] - \left[\frac{1}{2} (1)^2 \ln 1 - \frac{1}{4} (1)^2 \right]$$

$$8 \ln 4 - 4 + \frac{1}{4}$$

$$8 \ln 4 - \frac{15}{4}$$

$$\frac{1}{4} (32 \ln 4 - 15)$$

$$\boxed{\ln 4 = 2 \ln 2}$$

$$\frac{1}{4} (64 \ln 2 - 15)$$



3. The curve C has the equation

$$\cos 2x + \cos 3y = 1, \quad -\frac{\pi}{4} \leq x \leq \frac{\pi}{4}, \quad 0 \leq y \leq \frac{\pi}{6}$$

(a) Find $\frac{dy}{dx}$ in terms of x and y . (3)

The point P lies on C where $x = \frac{\pi}{6}$.

(b) Find the value of y at P . (3)

(c) Find the equation of the tangent to C at P , giving your answer in the form $ax + by + c\pi = 0$, where a , b and c are integers. (3)

$$-2 \sin 2x - 3 \sin 3y \frac{dy}{dx} = 0$$

$$-3 \sin 3y \frac{dy}{dx} = 2 \sin 2x$$

$$\frac{dy}{dx} = -\frac{2 \sin 2x}{3 \sin 3y}$$

b/ $x = \frac{\pi}{6}$

$$\cos\left(\frac{2\pi}{6}\right) + \cos 3y = 1$$

$$\cos 3y = \frac{1}{2}$$

$$3y = \frac{1}{3}\pi,$$

$$y = \underline{\underline{\frac{1}{9}\pi}}$$

c/ $\frac{dy}{dx} = \frac{-2 \sin\left(\frac{2\pi}{6}\right)}{3 \sin\left(\frac{3\pi}{9}\right)}$

$$= \frac{-2\sqrt{3}}{3} = -\frac{2}{3}$$

$$y = mx + c$$



Question 3 continued

$$\frac{1}{9}\pi = -\frac{2}{3} \cdot \frac{\pi}{6} + c$$

$$\frac{1}{9}\pi = -\frac{1}{9}\pi + c$$

$$c = \frac{2}{9}\pi$$

$$y = -\frac{2}{3}x + \frac{2}{9}\pi$$

$$9y = -6x + 2\pi$$

$$\underline{\underline{6x + 9y - 2\pi = 0}}$$



4. The line l_1 has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix}$$

and the line l_2 has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -4 \\ 1 \end{pmatrix}$$

where λ and μ are parameters.

The lines l_1 and l_2 intersect at the point A and the acute angle between l_1 and l_2 is θ .

(a) Write down the coordinates of A . (1)

(b) Find the value of $\cos \theta$. (3)

The point X lies on l_1 where $\lambda = 4$.

(c) Find the coordinates of X . (1)

(d) Find the vector \overrightarrow{AX} . (2)

(e) Hence, or otherwise, show that $|\overrightarrow{AX}| = 4\sqrt{26}$. (2)

The point Y lies on l_2 . Given that the vector \overrightarrow{YX} is perpendicular to l_1 ,

(f) find the length of AY , giving your answer to 3 significant figures. (3)

a/ $\begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix}$

b/ $\cos \theta = \frac{a \cdot b}{|a||b|}$

$a \cdot b = 4(3) + -1(-4) + 3(1)$
 $= 19$



Question 4 continued

$$|a| = \sqrt{4^2 + 1^2 + 3^2}$$

$$= \sqrt{26}$$

$$|b| = \sqrt{3^2 + 4^2 + 1^2}$$

$$= \sqrt{26}$$

$$\cos \theta = \frac{19}{\sqrt{26}\sqrt{26}}$$

~~$$= \frac{19}{26}$$~~

$$\cos \theta = \frac{19}{26}$$

c) $\lambda = 4$


$$\begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + 4 \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix} = \begin{pmatrix} 10 \\ 0 \\ 11 \end{pmatrix}$$

d)

$$\begin{pmatrix} 10 \\ 0 \\ 11 \end{pmatrix} - \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} = \begin{pmatrix} 16 \\ -4 \\ 12 \end{pmatrix}$$

e) $\sqrt{16^2 + 4^2 + 12^2} = \sqrt{416} = \underline{\underline{4\sqrt{26}}}$

f)



~~$$\tan \theta = \frac{x}{4\sqrt{26}}$$~~

~~$$x = 4\sqrt{26} \tan \theta$$~~

$$\cos \theta = \frac{4\sqrt{26}}{x}$$

$$x = \frac{4\sqrt{26}}{19/26}$$

$$= 27.9 \quad (3 \text{ s.f.})$$



5. (a) Find $\int \frac{9x+6}{x} dx, x > 0.$

(2)

(b) Given that $y = 8$ at $x = 1$, solve the differential equation

$$\frac{dy}{dx} = \frac{(9x+6)y^{\frac{1}{3}}}{x}$$

giving your answer in the form $y^2 = g(x).$

(6)

a) $\int 9 + \frac{6}{x} dx$

$$= 9x + 6 \ln x + c$$

b) $\int y^{-1/3} dy = \int \frac{9x+6}{x} dx$

$$\frac{3}{2} y^{2/3} = 9x + 6 \ln x + c$$

(1, 8)
~~(8, 1)~~

~~$$\frac{3}{2} (1)^{2/3} = 9(8) + 6 \ln(8) + c$$~~

~~$$\frac{3}{2} = 72 + 6 \ln 8$$~~

$$\frac{3}{2} (8)^{2/3} = 9(1) + 6 \ln 1 + c$$

$$6 = 9 + c$$

$$c = -3$$

$$\frac{3}{2} y^{2/3} = 9x + 6 \ln x - 3$$

$$y^{2/3} = 6x + 4 \ln x - 2$$

$$y^2 = (6x + 4 \ln x - 2)^3$$



6. The area A of a circle is increasing at a constant rate of $1.5 \text{ cm}^2 \text{ s}^{-1}$. Find, to 3 significant figures, the rate at which the radius r of the circle is increasing when the area of the circle is 2 cm^2 .

(5)

$$\frac{dA}{dt} = 1.5$$

$$\frac{dr}{dt} = \frac{dA}{dt} \times \frac{dr}{dA}$$

$$= 1.5 \times \frac{1}{2\pi r}$$

$$= \frac{1.5}{2\pi r}$$

$$\frac{dr}{dA}$$

$$A = \pi r^2$$

~~2d~~

$$\frac{dA}{dr} = 2\pi r$$

$$\frac{dr}{dA} = \frac{1}{2\pi r}$$

when $A = 2$ $2 = \pi r^2$

$$r = \sqrt{\frac{2}{\pi}}$$

$$\frac{dr}{dt} = \frac{1.5}{2\pi \left(\sqrt{\frac{2}{\pi}}\right)} = \underline{\underline{0.299}} \text{ s}^{-1}$$



7.




Figure 2

Figure 2 shows a sketch of the curve C with parametric equations

$$x = 5t^2 - 4, \quad y = t(9 - t^2)$$

The curve C cuts the x -axis at the points A and B .

- (a) Find the x -coordinate at the point A and the x -coordinate at the point B . (3)

The region R , as shown shaded in Figure 2, is enclosed by the loop of the curve.

- (b) Use integration to find the area of R . (6)

a) A and B : $y = 0$

$$\begin{aligned} 0 &= t(9 - t^2)^2 \\ &= t(3 - t)(3 + t) \\ t &= 0 \quad t = 3 \quad t = -3 \end{aligned}$$

$$\begin{aligned} x &= 5(0)^2 - 4 \\ &= -4 \end{aligned}$$

$$\begin{aligned} x &= 5(3)^2 - 4 \\ &= 41 \end{aligned}$$

$$A : (-4, 0)$$

$$B : (41, 0)$$

b/ $\int_{-4}^{41} y \, dx$



Question 7 continued

$$\int_0^3 y \frac{dx}{dt} dt$$

$$\int_0^3 t(9-t^2)(10t) dt$$

$$\int_0^3 10t^2(9-t^2) dt$$

$$\int_0^3 90t^2 - 10t^4 dt$$

$$\left[\frac{90t^3}{3} - \frac{10t^5}{5} + C \right]_0^3$$

$$\left[30t^3 - 2t^5 \right]_0^3$$

$$30(3)^3 - 2(3)^5 - 0$$

$$= 324 \text{ units}^2$$

$$324 \times 2 = \underline{\underline{648 \text{ units}^2}}$$



8. (a) Using the substitution $x = 2 \cos u$, or otherwise, find the exact value of

$$\int_1^{\sqrt{2}} \frac{1}{x^2 \sqrt{4-x^2}} dx \tag{7}$$




Figure 3

Figure 3 shows a sketch of part of the curve with equation $y = \frac{4}{x(4-x^2)^{\frac{1}{2}}}$, $0 < x < 2$.

The shaded region S , shown in Figure 3, is bounded by the curve, the x -axis and the lines with equations $x = 1$ and $x = \sqrt{2}$. The shaded region S is rotated through 2π radians about the x -axis to form a solid of revolution.

(b) Using your answer to part (a), find the exact volume of the solid of revolution formed.

(3)

a/

$$\begin{aligned} \sqrt{2} &= 2 \cos u & 1 &= 2 \cos u & \frac{dx}{du} &= -2 \sin u \\ u &= \frac{1}{4} \pi & u &= \frac{1}{3} \pi & & \end{aligned}$$

$$\int_{\frac{1}{3}\pi}^{\frac{1}{4}\pi} \frac{1}{(2 \cos u)^2 \sqrt{4 - (2 \cos u)^2}} \frac{dx}{du} du$$

$$\int_{\frac{1}{3}\pi}^{\frac{1}{4}\pi} \frac{1}{4 \cos^2 u \sqrt{4 - 4 \cos^2 u}} \cdot -2 \sin u du$$

$$\int_{\frac{1}{3}\pi}^{\frac{1}{4}\pi} \frac{-2 \sin u}{4 \cos^2 u \sqrt{4 \sin^2 u}} du$$



Question 8 continued

$$\int_{\frac{1}{3}\pi}^{\frac{1}{4}\pi} \frac{-2 \sin u}{4 \cos^2 u (2 \sin u)} du$$

$$\int_{\frac{1}{3}\pi}^{\frac{1}{4}\pi} \frac{-1}{4 \cos^2 u} du$$

$$\frac{1}{4} \int_{\frac{1}{4}\pi}^{\frac{1}{3}\pi} \sec^2 u du$$

~~$$\frac{1}{4} \int_{\frac{1}{4}}^{\frac{1}{3}\pi}$$~~

$$\frac{1}{4} \left[\tan u + c \right]_{\frac{1}{4}\pi}^{\frac{1}{3}\pi}$$

$$\frac{1}{4} \left[(\sqrt{3}) - (1) \right]$$

$$\frac{\sqrt{3} - 1}{4}$$

b)

$$\pi \int_1^{\sqrt{2}} \left(\frac{4}{x(4-x^2)^{1/4}} \right)^2 dx$$

$$\pi \int_1^{\sqrt{2}} \frac{16}{x^2 \sqrt{(4-x)^2}} dx$$

$$16\pi \int_1^{\sqrt{2}} \frac{1}{x^2 \sqrt{(4-x)^2}} dx$$

$$4 \times 16\pi \left(\frac{\sqrt{3} - 1}{4} \right) = \underline{\underline{4\pi(\sqrt{3} - 1)}}$$

Q8

(Total 10 marks)

TOTAL FOR PAPER: 75 MARKS

END

