



1. A curve  $C$  has the equation

$$x^3 + 2xy - x - y^3 - 20 = 0$$

(a) Find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

$$u = 2x \quad v = y$$

$$\frac{dy}{dx} = 2 \frac{dv}{dx} = \frac{dy}{dx}$$

(5)

(b) Find an equation of the tangent to  $C$  at the point  $(3, -2)$ , giving your answer in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers.

(2)

$$a) \quad 3x^2 + 2y + 2x \frac{dy}{dx} - 1 - 3y^2 \frac{dy}{dx} = 0$$

$$3x^2 + 2y - 1 = 3y^2 \frac{dy}{dx} - 2x \frac{dy}{dx}$$

$$3x^2 + 2y - 1 = \frac{dy}{dx} (3y^2 - 2x)$$

$$\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{3y^2 - 2x}$$

$$b) \quad \begin{matrix} x & y \\ (3, & -2) \end{matrix}$$

$$\frac{dy}{dx} = \frac{3(3)^2 + 2(-2) - 1}{3(-2)^2 - 2(3)}$$

$$= \frac{11}{3}$$

$$y = \frac{11}{3}x + c$$

$$-2 = \frac{11}{3}(3) + c$$

$$-2 = 11 + c$$

$$c = -13$$

$$y = \frac{11}{3}x - 13$$

$$3y = 11x - 39$$

$$\underline{11x - 3y - 39 = 0}$$



2. Given that the binomial expansion of  $(1 + kx)^{-4}$ ,  $|kx| < 1$ , is

$$1 - 6x + Ax^2 + \dots$$

(a) find the value of the constant  $k$ , (2)

(b) find the value of the constant  $A$ , giving your answer in its simplest form. (3)

$$1 + (-4)(kx) + \frac{(-4)(-5)}{2}(kx)^2$$

a/

$$-4k = -6$$

$$4k = 6$$

$$k = \frac{3}{2}$$

b/

$$10k^2 = A$$

$$10\left(\frac{3}{2}\right)^2 = A$$

$$10\left(\frac{9}{4}\right) = A$$

$$A = \frac{90}{4}$$

$$= \frac{45}{2}$$



3.




Figure 1

Figure 1 shows a sketch of part of the curve with equation  $y = \frac{10}{2x + 5\sqrt{x}}$ ,  $x > 0$

The finite region  $R$ , shown shaded in Figure 1, is bounded by the curve, the  $x$ -axis, and the lines with equations  $x = 1$  and  $x = 4$

The table below shows corresponding values of  $x$  and  $y$  for  $y = \frac{10}{2x + 5\sqrt{x}}$

$x$	1	2	3	4
$y$	1.42857	0.90326	0.68212	0.55556

- Complete the table above by giving the missing value of  $y$  to 5 decimal places. (1)
- Use the trapezium rule, with all the values of  $y$  in the completed table, to find an estimate for the area of  $R$ , giving your answer to 4 decimal places. (3)
- By reference to the curve in Figure 1, state, giving a reason, whether your estimate in part (b) is an overestimate or an underestimate for the area of  $R$ . (1)
- Use the substitution  $u = \sqrt{x}$ , or otherwise, to find the exact value of

$$\int_1^4 \frac{10}{2x + 5\sqrt{x}} dx \quad (6)$$

$$\begin{aligned} & \text{b/ } \frac{1}{2} \left( \frac{1.42857}{2} + 0.90326 + 0.68212 + \frac{0.55556}{2} \right) \\ & = 2.5774 \text{ units}^2 \end{aligned}$$



Question 3 continued

c/ overestimate  
it is curving down.

$$d/ \int_1^4 \frac{10}{2x + 5\sqrt{x}} dx$$

$$u = x^{1/2}$$

$$\frac{du}{dx} = \frac{1}{2} x^{-1/2} \quad \frac{dx}{du} = 2x^{1/2}$$

$$x=4 \quad u=2$$

$$x=1 \quad u=1$$

$$\int_1^2 \frac{10}{2x + 5\sqrt{x}} \frac{dx}{du} du$$

$$\int_1^2 \frac{10}{2x + 5u} \cdot 2x^{1/2} du$$

$$\int_1^2 \frac{10}{2u^2 + 5u} \cdot 2u du$$

$$\int_1^2 \frac{20}{2u + 5} du$$

$$\left[ 10 \ln(2u + 5) \right]_1^2$$

$$10 \ln 9 - 10 \ln 7$$

$$10 \ln \left( \frac{9}{7} \right)$$



4.




Figure 2

A vase with a circular cross-section is shown in Figure 2. Water is flowing into the vase.

When the depth of the water is  $h$  cm, the volume of water  $V$  cm<sup>3</sup> is given by

$$V = 4\pi h(h + 4), \quad 0 \leq h \leq 25$$

Water flows into the vase at a constant rate of  $80\pi$  cm<sup>3</sup>s<sup>-1</sup>

Find the rate of change of the depth of the water, in cm s<sup>-1</sup>, when  $h = 6$

(5)

$$\frac{dV}{dt} = 80\pi$$

$$V = 4\pi h^2 + 16\pi h$$

$$\frac{dV}{dh} = 8\pi h + 16\pi$$

$$\frac{dh}{dt} = \frac{dV}{dt} \times \frac{dh}{dV}$$

$$= 80\pi \times \frac{1}{8\pi h + 16\pi}$$

$$= \frac{80\pi}{8\pi h + 16\pi} = \frac{10}{h + 2}$$

$$(h=6) \quad = \frac{10}{8} = \frac{5}{4} = \underline{\underline{1.25 \text{ cm s}^{-1}}}$$



5.




Figure 3

Figure 3 shows a sketch of the curve  $C$  with parametric equations

$$x = 4 \cos\left(t + \frac{\pi}{6}\right), \quad y = 2 \sin t, \quad 0 \leq t < 2\pi$$

(a) Show that

$$x + y = 2\sqrt{3} \cos t \tag{3}$$

(b) Show that a cartesian equation of  $C$  is

$$(x + y)^2 + ay^2 = b$$

where  $a$  and  $b$  are integers to be determined.

(2)

$$\begin{aligned} \text{a) } x + y &= 4 \cos\left(t + \frac{\pi}{6}\right) + 2 \sin t \\ &= 4\left(\cos t \cos \frac{\pi}{6} - \sin t \sin \frac{\pi}{6}\right) + 2 \sin t \\ &= 4\left(\frac{\sqrt{3}}{2} \cos t - \frac{1}{2} \sin t\right) + 2 \sin t \\ &= 2\sqrt{3} \cos t - 2 \sin t + 2 \sin t \\ &= \underline{2\sqrt{3} \cos t} \end{aligned}$$

$$\begin{aligned} \text{b) } (x + y)^2 &= 12 \cos^2 t \\ (x + y)^2 &= 12 - 12 \sin^2 t \end{aligned}$$

$$\boxed{y^2 = 4 \sin^2 t}$$





Question 5 continued

$$3y^2 = 12 \sin^2 t$$

$$(x+y)^2 = 12 - 3y^2$$

$$\underline{\underline{(x+y)^2 + 3y^2 = 12}}$$

Q5

(Total 11 marks)





6. (i) Find

$$\int x e^{4x} dx \quad (3)$$

(ii) Find

$$\int \frac{8}{(2x-1)^3} dx, \quad x > \frac{1}{2} \quad (2)$$

(iii) Given that  $y = \frac{\pi}{6}$  at  $x = 0$ , solve the differential equation

$$\frac{dy}{dx} = e^x \operatorname{cosec} 2y \operatorname{cosec} y \quad (7)$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

$$u = x \quad \frac{dv}{dx} = e^{4x}$$

$$\frac{du}{dx} = 1 \quad v = \frac{1}{4} e^{4x}$$

$$= \frac{1}{4} x e^{4x} - \int \frac{1}{4} e^{4x} dx$$

$$= \frac{1}{4} x e^{4x} - \frac{1}{16} e^{4x} + C$$

ii)  $\int 8(2x-1)^{-3} dx$

$$= \underline{\underline{-2(2x-1)^{-2}}}$$

iii)  $\int \frac{1}{\operatorname{cosec} 2y \operatorname{cosec} y} dy = \int e^x dx$

$$\int \sin 2y \sin y dy = e^x + C$$

$$\int \frac{\cancel{2 \sin^2 y} + 1}{2 \sin y \cos y} (\sin y) dy = e^{2x} + C$$



## Question 6 continued

$$\int 2 \sin^2 y \cos y \, dy = e^x + c$$

$$\frac{2}{3} \sin^3 y = e^x + c$$

$$\frac{2}{3} \sin^3\left(\frac{\pi}{6}\right) = e^a + c$$

$$\begin{matrix} x & y \\ \hline 0 & \frac{\pi}{6} \end{matrix}$$

$$\frac{1}{12} = 1 + c$$

$$c = -\frac{11}{12}$$

$$\frac{2}{3} \sin^3 y = e^x - \frac{11}{12}$$

Q6

(Total ~~11~~ marks)

21

Turn over



P 4 1 8 2 8 A 0 2 1 2 8

7.




Figure 4

Figure 4 shows a sketch of part of the curve  $C$  with parametric equations

$$x = 3 \tan \theta, \quad y = 4 \cos^2 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point  $P$  lies on  $C$  and has coordinates  $(3, 2)$ .

The line  $l$  is the normal to  $C$  at  $P$ . The normal cuts the  $x$ -axis at the point  $Q$ .

- (a) Find the  $x$  coordinate of the point  $Q$ . (6)

The finite region  $S$ , shown shaded in Figure 4, is bounded by the curve  $C$ , the  $x$ -axis, the  $y$ -axis and the line  $l$ . This shaded region is rotated  $2\pi$  radians about the  $x$ -axis to form a solid of revolution.

- (b) Find the exact value of the volume of the solid of revolution, giving your answer in the form  $p\pi + q\pi^2$ , where  $p$  and  $q$  are rational numbers to be determined.

[You may use the formula  $V = \frac{1}{3}\pi r^2 h$  for the volume of a cone.] (9)

$$a) \quad \frac{dx}{d\theta} = 3 \sec^2 \theta \quad \frac{dy}{d\theta} = -8 \cos \theta \sin \theta$$

$$\frac{dy}{dx} = \frac{-8 \cos \theta \sin \theta}{3 \sec^2 \theta}$$

$$= -\frac{8}{3} \cos^3 \theta \sin \theta$$

$$\theta = \frac{1}{4} \pi \quad [3 = 3 \tan \theta]$$



Question 7 continued

$$\frac{dy}{dx} = -\frac{8}{3} \left(\cos \frac{1}{4}\pi\right)^3 \sin \frac{1}{4}\pi$$

$$= -\frac{2}{3}$$

$$\therefore m = \frac{3}{2}$$

$$y = \frac{3}{2}x + c \quad (3, 2)$$

$$2 = \frac{3}{2}(3) + c$$

$$2 = \frac{9}{2} + c$$

$$c = -\frac{5}{2}$$

$$y = \frac{3}{2}x - \frac{5}{2}$$

crosses x when  $y=0$

$$0 = \frac{3}{2}x - \frac{5}{2}$$

$$5 = 3x$$

$$x = \frac{5}{3}$$

$$\pi \int_0^3 y^2 dx$$

$$\pi \int_0^3 (4 \cos^2 \theta)^2 dx$$

$$x=3 \quad \theta = \frac{1}{4}\pi$$

$$x=0 \quad \theta = 0$$

$$\pi \int_0^{\frac{\pi}{4}} 16 \cos^4 \theta \frac{dx}{d\theta} d\theta$$

$$\pi \int_0^{\frac{\pi}{4}} 48 \cos^2 \theta d\theta$$

$$\frac{dx}{d\theta} = \frac{3 \cos^2 \theta}{\cos^2 \theta}$$



7  
Question 4 continued

$$48\pi \int_0^{\pi/4} \cos^2 \theta \, d\theta$$

$$48\pi \int_0^{\pi/4} \frac{1}{2} \cos 2\theta + \frac{1}{2} \, d\theta$$

$\cos 2\theta = 2\cos^2 \theta - 1$   
 $\frac{1}{2} \cos 2\theta + \frac{1}{2} = \cos^2 \theta$

$$48\pi \left[ \frac{1}{4} \sin 2\theta + \frac{1}{2} \theta \right]_0^{\pi/4}$$

$$48\pi \left[ \frac{1}{4} + \frac{\pi}{8} \right]$$

$$\underline{\underline{12\pi + 6\pi^2}}$$

Cone:  $h = \frac{4}{3}$   
 $r = 2$

$$V = \frac{1}{3} \pi (2)^2 \left(\frac{4}{3}\right)$$

$$= \frac{16}{9} \pi$$

$$\underline{\underline{\frac{92}{9} \pi + 6\pi^2}}$$



8. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $\begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix}$

and the point  $B$  has position vector  $\begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix}$

The line  $l_1$  passes through the points  $A$  and  $B$ .

(a) Find the vector  $\vec{AB}$ . (2)

(b) Hence find a vector equation for the line  $l_1$ . (1)

The point  $P$  has position vector  $\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix}$

Given that angle  $PBA$  is  $\theta$ ,

(c) show that  $\cos \theta = \frac{1}{3}$ . (3)

The line  $l_2$  passes through the point  $P$  and is parallel to the line  $l_1$

(d) Find a vector equation for the line  $l_2$ . (2)

The points  $C$  and  $D$  both lie on the line  $l_2$

Given that  $AB = PC = DP$  and the  $x$  coordinate of  $C$  is positive,

(e) find the coordinates of  $C$  and the coordinates of  $D$ . (3)

(f) find the exact area of the trapezium  $ABCD$ , giving your answer as a simplified surd. (4)

a/  $\vec{AB} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} - \begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$

b/  $r = \begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$



Question 8 continued

$$\vec{BA} = \begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}$$

$$\vec{BP} = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}$$

$$\cos \theta = \frac{a \cdot b}{|a| |b|}$$

$$a \cdot b = -1(1) + 1(-1) + -1(-5)$$

$$= -1 - 1 + 5$$

$$= 3$$

$$|a| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$$

$$|b| = \sqrt{1^2 + 1^2 + 5^2} = \sqrt{27}$$


$$\cos \theta = \frac{3}{\sqrt{3}\sqrt{27}}$$

$$= \frac{3}{\sqrt{81}} = \frac{3}{9} = \underline{\underline{\frac{1}{3}}}$$

d/

$$r = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$$

e/



$$\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ 4 \end{pmatrix}$$






Question 8 continued



$$C: \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix} \quad D: \begin{pmatrix} -1 \\ 3 \\ 4 \end{pmatrix}$$

f/ distance  $\vec{BP} = \sqrt{1^2 + 1^2 + 5^2}$   
 $= \sqrt{27}$



$$\sin \theta = \frac{h}{\sqrt{27}}$$

$$h = 2\sqrt{6}$$



$$\text{Area} = \frac{2\sqrt{3} + \sqrt{3}}{2} \cdot 2\sqrt{6}$$

$$= \frac{3\sqrt{3}}{2} \cdot 2\sqrt{6}$$

$$= \underline{\underline{9\sqrt{2}}}$$

