


1. Two particles B and C have mass m kg and 3 kg respectively. They are moving towards each other in opposite directions on a smooth horizontal table. The two particles collide directly. Immediately before the collision, the speed of B is 4 m s^{-1} and the speed of C is 2 m s^{-1} . In the collision the direction of motion of C is reversed and the direction of motion of B is unchanged. Immediately after the collision, the speed of B is 1 m s^{-1} and the speed of C is 3 m s^{-1} .

Find

- (a) the value of m , (3)

- (b) the magnitude of the impulse received by C . (2)



$$a) \quad m_1 u_1 + m_2 u_2 = m_1 v_1 + m_2 v_2$$

$$m(4) + 3(-2) = m(1) + 3(3)$$

$$4m - 6 = m + 9$$

$$3m = 15$$

$$\underline{m = 5}$$

$$b/ \quad I = mv - mu$$

$$= 3(3) - 3(-2)$$

$$= \underline{15 \text{ N s}}$$



2. A ball is thrown vertically upwards with speed $u \text{ m s}^{-1}$ from a point P at height h metres above the ground. The ball hits the ground 0.75 s later. The speed of the ball immediately before it hits the ground is 6.45 m s^{-1} . The ball is modelled as a particle.

(a) Show that $u = 0.9$ (3)

(b) Find the height above P to which the ball rises before it starts to fall towards the ground again. (2)

(c) Find the value of h . (3)

$s =$

$u = ?$

$v = -6.45$

$a = -9.8$

$t = 0.75$

a/ $v = u + at$

$-6.45 = u - 7.35$

$u = \underline{0.9 \text{ ms}^{-1}}$

b/ $s =$

$u = 0.9$

$v^2 = u^2 + 2as$

$v = 0$

$0 = (0.9)^2 + 2(-9.8)s$

$a = -9.8$

~~$s = 3.97 \text{ m}$~~ (3st)

$t = ?$

$s = 0.0413 \text{ m}$

c/ whole motion

$s =$

$u = 0.9$

$s = ut + \frac{1}{2}at^2$

$v = -6.45$

$= (0.9)(0.75) + \frac{1}{2}(-9.8)(0.75)^2$

$a = -9.8$


$= -2.08 \text{ m}$ 3st.

$t = 0.75$

$\therefore \underline{h = 2.08 \text{ m}}$ 3st



3.



A uniform beam AB has mass 20 kg and length 6 m . The beam rests in equilibrium in a horizontal position on two smooth supports. One support is at C , where $AC = 1\text{ m}$, and the other is at the end B , as shown in Figure 1. The beam is modelled as a rod.

(a) Find the magnitudes of the reactions on the beam at B and at C . (5)

A boy of mass 30 kg stands on the beam at the point D . The beam remains in equilibrium. The magnitudes of the reactions on the beam at B and at C are now equal. The boy is modelled as a particle.

(b) Find the distance AD . (5)

a/ Taking moments about B:

$$5(R_c) = 60g$$

$$R_c = \underline{12g}$$

Forces up = forces down

$$R_c + R_B = 20g$$

$$12g + R_B = 20g$$

$$R_B = \underline{8g}$$

b/ forces up = forces down

$$2R = 50g$$

$$R = \underline{25g}$$

Taking moments about A:

$$30g x + 60g = 25g + 150g$$

$$30x = 115$$

$$x = \underline{\underline{\frac{23}{6}\text{ m}}}$$



4. A particle P of mass 2 kg is moving under the action of a constant force \mathbf{F} newtons. The velocity of P is $(2\mathbf{i} - 5\mathbf{j}) \text{ m s}^{-1}$ at time $t = 0$, and $(7\mathbf{i} + 10\mathbf{j}) \text{ m s}^{-1}$ at time $t = 5 \text{ s}$.

Find

- (a) the speed of P at $t = 0$, (2)
- (b) the vector \mathbf{F} in the form $a\mathbf{i} + b\mathbf{j}$, (5)
- (c) the value of t when P is moving parallel to \mathbf{i} . (4)

$$\begin{aligned} \text{a/ speed} &= \sqrt{2^2 + 5^2} \\ &= \sqrt{29} \text{ ms}^{-1} \end{aligned}$$

$$\begin{aligned} \text{b/ } \mathbf{a} &= \frac{7\mathbf{i} + 10\mathbf{j} - (2\mathbf{i} - 5\mathbf{j})}{5} \\ &= \frac{5\mathbf{i} + 15\mathbf{j}}{5} \\ &= \mathbf{i} + 3\mathbf{j} \text{ ms}^{-2} \end{aligned}$$

$$\begin{aligned} \mathbf{F} &= m\mathbf{a} \\ &= 2(\mathbf{i} + 3\mathbf{j}) \\ &= \underline{\underline{2\mathbf{i} + 6\mathbf{j}}} \end{aligned}$$

$$\text{c/ parallel to } \mathbf{i} \rightarrow \mathbf{j} = 0$$

$$\begin{aligned} \mathbf{v} &= \mathbf{v}_0 + \mathbf{a}t \\ &= (2\mathbf{i} - 5\mathbf{j}) + t(\mathbf{i} + 3\mathbf{j}) \end{aligned}$$

$$\begin{aligned} \text{ii/ } -5 + 3t &= 0 \\ 3t &= 5 \\ t &= \frac{5}{3} \text{ seconds} \end{aligned}$$



5. A car accelerates uniformly from rest for 20 seconds. It moves at constant speed $v \text{ m s}^{-1}$ for the next 40 seconds and then decelerates uniformly for 10 seconds until it comes to rest.

(a) For the motion of the car, sketch

(i) a speed-time graph,



(ii) an acceleration-time graph.

(6)

Given that the total distance moved by the car is 880 m,

(b) find the value of v .

(4)



Question 5 continued

$$b/ \left(\frac{40 + 76}{2} \right) v = 880$$

$$55v = 880$$

$$\underline{\underline{v = 16}}$$



6.




Figure 2

A particle of weight 120 N is placed on a fixed rough plane which is inclined at an angle α to the horizontal, where $\tan \alpha = \frac{3}{4}$.

The coefficient of friction between the particle and the plane is $\frac{1}{2}$.

The particle is held at rest in equilibrium by a horizontal force of magnitude 30 N, which acts in the vertical plane containing the line of greatest slope of the plane through the particle, as shown in Figure 2.

- (a) Show that the normal reaction between the particle and the plane has magnitude 114 N. (4)




Figure 3

The horizontal force is removed and replaced by a force of magnitude P newtons acting up the slope along the line of greatest slope of the plane through the particle, as shown in Figure 3. The particle remains in equilibrium.

- (b) Find the greatest possible value of P . (8)
- (c) Find the magnitude and direction of the frictional force acting on the particle when $P = 30$. (3)




Question 6 continued

$$\tan \alpha = \frac{3}{4}$$

$$\cos \alpha = \frac{4}{5}$$

$$\sin \alpha = \frac{3}{5}$$



$$\mu = \frac{1}{2}$$

a/ Resolving perp. to plane

$$R = 120 \cos \alpha + 30 \sin \alpha$$

$$= 120 \left(\frac{4}{5} \right) + 30 \left(\frac{3}{5} \right)$$

$$= \underline{\underline{114 \text{ N}}}$$

b/ $R = 120 \cos \alpha$

$$= 96 \text{ N}$$

Resolving parallel to plane:

$$P = 120 \sin \alpha + 0.5(96)$$


$$= 72 + 48$$

$$= \underline{\underline{120 \text{ N}}}$$

c/ Resolving parallel to plane.

$$30 + = 120 \sin \alpha + F$$

$$F = -42$$

42 N acting up the plane 



7.




Figure 4

Two particles A and B , of mass 7 kg and 3 kg respectively, are attached to the ends of a light inextensible string. Initially B is held at rest on a rough fixed plane inclined at angle θ to the horizontal, where $\tan \theta = \frac{5}{12}$. The part of the string from B to P is parallel to a line of greatest slope of the plane. The string passes over a small smooth pulley, P , fixed at the top of the plane. The particle A hangs freely below P , as shown in Figure 4. The coefficient of friction between B and the plane is $\frac{2}{3}$. The particles are released from rest with the string taut and B moves up the plane.

(a) Find the magnitude of the acceleration of B immediately after release. (10)

(b) Find the speed of B when it has moved 1 m up the plane. (2)

When B has moved 1 m up the plane the string breaks. Given that in the subsequent motion B does not reach P ,

(c) find the time between the instants when the string breaks and when B comes to instantaneous rest. (4)

$$\tan \theta = \frac{5}{12}$$

$$\cos \theta = \frac{12}{13}$$

$$\sin \theta = \frac{5}{13}$$

$$F = ma$$

$$A: 7g - T = 7a \quad (1)$$

B : Resolving perp. to plane:

$$R = 3g \cos \theta = \frac{36}{13}g$$



Question 7 continued

b) $F = ma$

$$T - 3g \sin \theta - \frac{24}{13}g = 3a$$

$$T - \frac{15}{13}g - \frac{24}{13}g = 3a$$

$$\begin{aligned} T - 3g &= 3a & (2) \\ -T + 7g &= 7a & (1) \end{aligned}$$

$$4g = 10a$$

$$\underline{\underline{a = 0.4g \text{ ms}^{-2}}}$$

b) $s = 1$

$u = 0$

$v = ?$

$a = 0.4g$

$t =$

$$v^2 = u^2 + 2as$$

$$v^2 = 2(0.4g)(1)$$

$$\underline{\underline{v = 2.8 \text{ ms}^{-1}}}$$

c) $F = ma$

$$-3g = 3a$$

$$a = -g$$

$s =$

$u = 2.8$

$v = 0$

$a = -g$

$t = ?$

$$v = u + at$$

$$0 = 2.8 + (-9.8)t$$

$$t = \frac{2}{9.8} \text{ seconds}$$

