




1. Each of 60 students was asked to draw a  $20^\circ$  angle without using a protractor. The size of each angle drawn was measured. The results are summarised in the box plot below.



- (a) Find the range for these data. (1)
- (b) Find the interquartile range for these data. (1)

The students were then asked to draw a  $70^\circ$  angle. The results are summarised in the table below.

Angle, $a$ , (degrees)	Number of students
$55 \leq a < 60$	6
$60 \leq a < 65$	15
$65 \leq a < 70$	13
$70 \leq a < 75$	11
$75 \leq a < 80$	8
$80 \leq a < 85$	7

- (c) Use linear interpolation to estimate the size of the median angle drawn. Give your answer to 1 decimal place. (2)
- (d) Show that the lower quartile is  $63^\circ$  (2)

For these data, the upper quartile is  $75^\circ$ , the minimum is  $55^\circ$  and the maximum is  $84^\circ$

An outlier is an observation that falls either more than  $1.5 \times$  (interquartile range) above the upper quartile or more than  $1.5 \times$  (interquartile range) below the lower quartile.

- (e) (i) Show that there are no outliers for these data.
- (ii) Draw a box plot for these data on the grid on page 3. (5)
- (f) State which angle the students were more accurate at drawing. Give reasons for your answer. (3)





Question 1 continued

Lined writing area with 20 horizontal lines.





2. An estate agent recorded the price per square metre,  $p$  £/m<sup>2</sup>, for 7 two-bedroom houses. He then coded the data using the coding  $q = \frac{p - a}{b}$ , where  $a$  and  $b$  are positive constants. His results are shown in the table below.

$p$	1840	1848	1830	1824	1819	1834	1850
$q$	4.0	4.8	3.0	2.4	1.9	3.4	5.0

- (a) Find the value of  $a$  and the value of  $b$  **(2)**

The estate agent also recorded the distance,  $d$  km, of each house from the nearest train station. The results are summarised below.

$$S_{dd} = 1.02 \quad S_{qq} = 8.22 \quad S_{dq} = -2.17$$

- (b) Calculate the product moment correlation coefficient between  $d$  and  $q$  **(2)**
- (c) Write down the value of the product moment correlation coefficient between  $d$  and  $p$  **(1)**

The estate agent records the price and size of 2 additional two-bedroom houses,  $H$  and  $J$ .

House	Price (£)	Size (m <sup>2</sup> )
$H$	156 400	85
$J$	172 900	95

- (d) Suggest which house is most likely to be closer to a train station. Justify your answer. **(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question 2 continued

Lined writing area for question response.



P 4 4 8 4 5 A 0 7 2 4





**Question 2 continued**

Handwriting practice lines for Question 2 continued.

**(Total 8 marks)**

**Q2**

Q2



3. A college has 80 students in Year 12.

- 20 students study Biology
- 28 students study Chemistry
- 30 students study Physics
- 7 students study both Biology and Chemistry
- 11 students study both Chemistry and Physics
- 5 students study both Physics and Biology
- 3 students study all 3 of these subjects

(a) Draw a Venn diagram to represent this information. **(5)**

A Year 12 student at the college is selected at random.

(b) Find the probability that the student studies Chemistry but not Biology or Physics. **(1)**

(c) Find the probability that the student studies Chemistry or Physics or both. **(2)**

Given that the student studies Chemistry or Physics or both,

(d) find the probability that the student does not study Biology. **(2)**

(e) Determine whether studying Biology and studying Chemistry are statistically independent. **(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









4. Statistical models can provide a cheap and quick way to describe a real world situation.

- (a) Give two other reasons why statistical models are used. (2)

A scientist wants to develop a model to describe the relationship between the average daily temperature,  $x$  °C, and her household’s daily energy consumption,  $y$  kWh, in winter.

A random sample of the average daily temperature and her household’s daily energy consumption are taken from 10 winter days and shown in the table.

$x$	-0.4	-0.2	0.3	0.8	1.1	1.4	1.8	2.1	2.5	2.6
$y$	28	30	26	25	26	27	26	24	22	21

[You may use  $\sum x^2 = 24.76$   $\sum y = 255$   $\sum xy = 283.8$   $S_{xx} = 10.36$ ]

- (b) Find  $S_{xy}$  for these data. (3)

- (c) Find the equation of the regression line of  $y$  on  $x$  in the form  $y = a + bx$

Give the value of  $a$  and the value of  $b$  to 3 significant figures. (4)

- (d) Give an interpretation of the value of  $a$  (1)

- (e) Estimate her household’s daily energy consumption when the average daily temperature is 2°C (2)

The scientist wants to use the linear regression model to predict her household’s energy consumption in the summer.

- (f) Discuss the reliability of using this model to predict her household’s energy consumption in the summer. (2)

---

---

---

---

---

---

---

---

---

---

---

---









Leave  
blank

**Question 4 continued**

Lined writing area for the answer to Question 4.

**(Total 14 marks)**

**Q4**

Grading box for Question 4, consisting of two adjacent empty rectangular boxes.



5. In a quiz, a team gains 10 points for every question it answers correctly and loses 5 points for every question it does not answer correctly. The probability of answering a question correctly is 0.6 for each question. One round of the quiz consists of 3 questions.

The discrete random variable  $X$  represents the total number of points scored in one round. The table shows the incomplete probability distribution of  $X$

$x$	30	15	0	-15
$P(X=x)$	0.216			0.064

- (a) Show that the probability of scoring 15 points in a round is 0.432 (2)
- (b) Find the probability of scoring 0 points in a round. (1)
- (c) Find the probability of scoring a total of 30 points in 2 rounds. (3)
- (d) Find  $E(X)$  (2)
- (e) Find  $\text{Var}(X)$  (3)

In a bonus round of 3 questions, a team gains 20 points for every question it answers correctly and loses 5 points for every question it does not answer correctly.

- (f) Find the expected number of points scored in the bonus round. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









6. The random variable  $Z \sim N(0, 1)$

$A$  is the event  $Z > 1.1$

$B$  is the event  $Z > -1.9$

$C$  is the event  $-1.5 < Z < 1.5$

(a) Find

(i)  $P(A)$

(ii)  $P(B)$

(iii)  $P(C)$

(iv)  $P(A \cup C)$

**(6)**

The random variable  $X$  has a normal distribution with mean 21 and standard deviation 5

(b) Find the value of  $w$  such that  $P(X > w \mid X > 28) = 0.625$

**(6)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





**Question 6 continued**

Lined area for writing the answer to Question 6.

**(Total 12 marks)**

**Q6**

--	--

**TOTAL FOR PAPER: 75 MARKS**

**END**

