

# Mark Scheme (Results)

November 2013

Pearson Edexcel GCSE  
in Mathematics Linear (1MA0)  
Higher (Non-Calculator) Paper 1H

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

November 2013

Publications Code UG037492

All the material in this publication is copyright

© Pearson Education Ltd 2013

## NOTES ON MARKING PRINCIPLES

- 1 All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- 2 Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- 3 All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- 4 Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- 5 Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- 6 Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
  - i) *ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear*  
Comprehension and meaning is clear by using correct notation and labeling conventions.
  - ii) *select and use a form and style of writing appropriate to purpose and to complex subject matter*  
Reasoning, explanation or argument is correct and appropriately structured to convey mathematical reasoning.
  - iii) *organise information clearly and coherently, using specialist vocabulary when appropriate.*  
The mathematical methods and processes used are coherently and clearly organised and the appropriate mathematical vocabulary used.

**7 With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

If there is no answer on the answer line then check the working for an obvious answer.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

**8 Follow through marks**

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

**9 Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

**10 Probability**

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

**11 Linear equations**

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

**12 Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

**13 Range of answers**

Unless otherwise stated, when an answer is given as a range (e.g 3.5 – 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

**Guidance on the use of codes within this mark scheme**

M1 – method mark  
A1 – accuracy mark  
B1 – Working mark  
C1 – communication mark  
QWC – quality of written communication  
oe – or equivalent  
cao – correct answer only  
ft – follow through  
sc – special case  
dep – dependent (on a previous mark or conclusion)  
indep – independent  
isw – ignore subsequent working

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
1			90 450 225 1.5 960	3	M1 for $6 \div 4 (= 1.5)$ or $4 \div 6 (= 0.66..)$ or $\div 4 \times 6$ oe or sight of any one of the correct answers A1 for three correct A1 for all correct
2	(a)  (b)  (c)		Plot (90,17)  Positive  In range 16 to 20	1  1  2	B1 cao  B1 Positive  M1 for a single straight line segment with positive gradient that could be used as a line of best fit or a vertical line from 110 or a point plotted at (110, y) where y is in the range 16 to 20 A1 for an answer in the range 16 to 20 inclusive
3			$120 \text{ cm}^3$	4	M1 for $\frac{1}{2} \times 3 \times 4$  M1 (dep) for ' $\frac{1}{2} \times 3 \times 4$ ' $\times 20$  A1 for 120 B1 (indep) for $\text{cm}^3$

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
4	(a)		$4y + 5x + 5$	2	M1 5x or 5 seen A1 cao
	(b)		$3x(3x - 2y)$	2	B2 for $3x(3x - 2y)$ (B1 for $x(9x - 6y)$ or $3(3x^2 - 2xy)$ or $3x(ax - by)$ where $a$ and $b$ are integers not equal to zero)
	(c)		$4x + 8$	1	B1 cao
	(d)		$x^2 - 2x - 15$	2	M1 for 4 terms correct with or without signs or 3 out of no more than 4 terms correct with correct signs A1 cao
5	(a)		0.25	1	B1 oe
	(b)		150	2	M1 for $0.75 \times 200$ oe A1 cao
6	(a)		Shape with vertices at $(-1, 3)$ , $(0, 6)$ , $(2, 6)$ , $(1, 3)$	1	B1 for correct shape in correct position
	(b)		Rotation centre $(0,0)$ $90^\circ$ anticlockwise	3	B1 rotation B1 (centre) $(0,0)$ B1 $90^\circ$ anticlockwise or $270^\circ$ clockwise Note: award no marks if more than one transformation is given

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
7	(i)	20, 40, 60 12, 24, 36, 48, 60	3 and 5 or any multiple of 3, 5	4	M1 attempts multiples of both 20 and 12 (at least 3 of each shown but condone errors if intention is clear) or identifies 60 or a multiple of 60 M1 (dep on M1) for a division by 20 or 12 or counts up 'multiples' or identifies a common multiple (implied if one answer is correct or answers reversed) A1 cheese slices (packets) 3, burgers (boxes) 5 or any multiple of 3, 5  OR  M1 for expansion of either 20 or 12 into factors M1 for demonstration that both expansions include 4 (or $2 \times 2$ ) A1 cao for cheese slices (packets) 3, burgers (boxes) 5
	(ii)	$20 = 4 \times 5 = 2 \times 2 \times 5$ $12 = 4 \times 3 = 2 \times 2 \times 3$	60		B1 for 60 or ft from their correct answer in (i) or ft 'common multiple'
8			38	5	M1 $3x - 5 = 19 - x$ M1 for a correct operation to collect the $x$ terms or the number terms on one side of an equation of the form $ax + b = cx + d$ A1 for $x = 6$ M1 for substituting their value of $x$ in the three expressions and adding <b>or</b> substituting their value of $x$ after adding the three expressions A1 cao



PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
9	(a)		Criticisms	2	B1 Qu 1 Overlapping boxes, no units B1 Qu 2 e.g. no time frame, non-specific responses, no number quantities, open to interpretation, no option for those who do not exercise
	(b)		Question given	2	B1 for a correct question with a time frame B1 for at least 3 correctly labelled non-overlapping response boxes (need not be exhaustive) or at least 3 response boxes that are exhaustive for all integer values of their time unit (could be overlapping) NB Units must be included in either question or response boxes to score full marks  [Do not allow inequalities in response boxes]

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
*10			Not enough, needs £133	5	<p>M1 for splitting the shape (or showing recognition of the “absent” rectangle) and using a correct method to find the area of one shape  M1 for a complete and correct method to find the total area  M1 for a complete method to find 70% of 19 (= 13.3) or 70% of their total cost or 70% of their area  A1 114(m<sup>2</sup>) and (£)133 <b>or</b> 114(m<sup>2</sup>) and (£)13.3(0) and 108(m<sup>2</sup>)  C1 (dep on M2) for a conclusion supported by their calculations</p> <p>OR</p> <p>M1 for a complete method for the number of tins required for one section of the area of the floor  M1 for a complete method to find the number of tins for the whole floor  M1 for a complete method to find 70% of their total number of tins and multiply by 19  A1 (£)133  C1 (dep on M2) for a conclusion supported by their calculations</p>
11			164	5	<p>M1 <math>200 \div (3+2)</math> (= 40) or an equivalent ratio seen  M1 (dep) <math>3 \times '40'</math> (= 120) or <math>2 \times '40'</math> (= 80) or 120: 80 or 80:120  M1 a complete method to find 70% of their total number of large letters e.g. <math>0.7 \times '80'</math> (=56)  M1 multiplies their three totals by the correct unit price and adds, e.g. <math>60(p) \times '120' + (£)1 \times '56' + (£)1.50 \times '24'</math>  A1 164</p>

PAPER: 1MA0\_1H

Question	Working	Answer	Mark	Notes												
12	<table border="1" data-bbox="454 308 817 394"> <tr> <td><math>x</math></td> <td>-2</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> </tr> <tr> <td><math>y</math></td> <td>-4</td> <td>-1</td> <td>2</td> <td>5</td> <td>8</td> </tr> </table>	$x$	-2	-1	0	1	2	$y$	-4	-1	2	5	8	$y = 3x + 2$ drawn	4	B1 for axes scaled and labelled  <b>(Table of values)</b> M1 for at least 2 correct attempts to find points by substituting values of $x$ M1 ft for plotting at least 2 of their points (any points from their table must be correctly plotted) A1 for correct line between $x = -2$ and $x = 2$  <b>(No table of values)</b> M1 for at least 2 correct points with no more than 2 incorrect points M1 for at least 2 correct points (and no incorrect points) plotted OR line segment of $y = 3x + 2$ drawn A1 for correct line between $x = -2$ and $x = 2$  <b>(Use of <math>y = mx + c</math>)</b> M1 for line drawn with gradient of 3 <b>OR</b> line drawn with $y$ intercept at 2 M1 for line drawn with gradient of 3 <b>AND</b> with $y$ intercept at 2 A1 for correct line between $x = -2$ and $x = 2$  SC B2 (indep of B1) for correct line segment between $x = 0$ and $x = 2$ (ignore any additional incorrect line segment(s))
$x$	-2	-1	0	1	2											
$y$	-4	-1	2	5	8											

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
13		$35 \times 10 = 350$ $33 \times 11 = 363$ $363 - 350 = 13$  OR $10 \times (35 - 33) = 20$ $33 - 20 = 13$	13	3	M1 $35 \times 10 (= 350)$ or $33 \times 11 (= 363)$ M1 (dep) finding the difference in their totals e.g. '363' – '350' A1 cao  OR M1 $10 \times (35 - 33) (=20)$ or $11 \times (35 - 33) (=22)$ M1 (dep) $33 - '20'$ or $35 - '22'$ A1 cao
14	(a)		$\frac{1}{5}$	1	B1 oe
	(b)		$\frac{1}{9}$	1	B1 cao
	(c)	$9 \times 10^4 \times 3 \times 10^3$	$2.7 \times 10^8$	2	M1 $27 \times 10^7$ oe or $9 \times 3 \times 10^{4+3}$ A1 cao

PAPER: 1MA0_1H				
Question	Working	Answer	Mark	Notes
15	$6x + 8y = 10$ $6x - 9y = 27$ $y = -1$ $3x - 4 = 5$ $3x = 9$ $x = 3$  OR  $9x + 12y = 15$ $8x - 12y = 36$ $x = 3$ $9 + 4y = 5$ $4y = -4$ $y = -1$	$x = 3, y = -1$	4	M1 for a correct process to eliminate either variable (condone one arithmetic error) A1 cao for either $x$ or $y$ M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao  OR  M1 for full method to rearrange and substitute to eliminate either variable (condone one arithmetic error) A1 cao for either $x$ or $y$ M1 (dep on M1) for correct substitution of found value into one of the equations or appropriate method after starting again (condone one arithmetic error) A1 cao  Trial and improvement scores 0 marks unless both $x$ and $y$ are correct
16	$120 \div 20 = 6$ $6^2 = 36$ $36 \times 300 = 10\,800$	10 800	3	M1 $120 \div 20 (= 6)$ oe, can be implied by $120^2 \div 20^2$ M1 ' $6^2 \times 300$ ' A1 cao
17	$(3,6,7)$ to $(-2,2,5)$ $(-5, -4, -2)$ $(-2- 5, 2 - 4, 5 - 2)$	$(-7, -2, 3)$	2	M1 for midpoint plus change or complete method for 2 out of 3 coordinates, can be implied by 2 correct values A1 cao

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
18	(a)		68	1	B1 cao
	* (b)	$120 \div 20 = 6$ $6^2 = 36$ $36 \times 300 = 10\,800$	Yes as $28 > 20$ or $35\% > 25\%$ or $53 < 60$	3	M1 for reading a value from graph at time = 60 (=28, accept 27 to 28) M1 for '28' $\div 80 \times 100$ (= 35) or $25 \div 100 \times 80$ (= 20) C1 (dep on M2) for correct decision based on their figures  OR  M1 for $25 \div 100 \times 80$ (= 20) M1 for reading a value from graph at cf = 20 (=53, accept 52 to 54) C1 (dep on M2) for correct decision based on their figures
	(c)	28, 53, 68, 76, 96	Box plot plotted	3	B1 for ends of whiskers at 28 and 96 with a box B1 ft for median at '68' inside a box B1 for ends of box at 53 (accept 52 to 54) and 76
19			0.82	3	M1 for $1 - 0.7$ (= 0.3) or $1 - 0.4$ (= 0.6) M1 for $1 - '0.3' \times '0.6'$ A1 for 0.82 oe  OR  M1 for $1 - 0.7$ (= 0.3) or $1 - 0.4$ (= 0.6) M1 $(0.7 \times 0.4) + (0.7 \times '0.6') + ('0.3' \times 0.4)$ A1 for 0.82 oe

PAPER: 1MA0_1H					
Question		Working	Answer	Mark	Notes
20	(a)		4	3	M1 for correct expansion to $32x - 8$ or multiplying both sides by $3x$ or dividing both sides by 4 M1 for a complete and correct method to isolate the $x$ terms and the number terms (condone one arithmetic error in multiplying out the bracket) A1 cao
	(b)	$\frac{2(y-6)-(y+3)}{(y+3)(y-6)}$	$\frac{y-15}{(y+3)(y-6)}$	3	M1 for common denominator of $(y+3)(y-6)$ M1 for $\frac{2(y-6)}{(y+3)(y-6)} - \frac{y+3}{(y+3)(y-6)}$ oe or $\frac{2(y-6)-(y+3)}{(y+3)(y-6)}$ oe A1 for $\frac{y-15}{(y+3)(y-6)}$ or $\frac{y-15}{y^2-3y-18}$
21			100	4	M1 $y = kx^2$ oe or $36 = k \times 3^2$ A1 $k = 4$ M1 (dep on M1) $(y =) '4' \times 5^2$ A1 cao

PAPER: 1MA0_1H					
Question	Working	Answer	Mark	Notes	
*22		$360 - y$		$180 - \frac{y}{2}$	<p>4</p> <p>M1 <math>ADC = \frac{y}{2}</math></p> <p>A1 <math>180 - \frac{y}{2}</math></p> <p>C2 (dep on M1) for both reasons  <u>Angle at centre is twice the angle at the circumference</u>  <u>Opposite angles in cyclic quadrilateral add to 180°</u>  (C1 (dep on M1) for one appropriate circle theorem reason)</p> <p>OR</p> <p>M1 reflex <math>AOC = 360 - y</math></p> <p>A1 <math>\frac{360 - y}{2}</math> oe</p> <p>C2 (dep on M1) for both reasons  <u>Angles around a point add up to 360°</u>  <u>Angle at centre is twice the angle at the circumference</u>  (C1 (dep on M1) for one appropriate circle theorem reason)</p>
23				<p>Triangle with vertices at  <math>(-1, -4)</math>,  <math>(-1, -5)</math>,  <math>(-3, -4.5)</math></p>	<p>2</p> <p>M1 for correct shape and size and the correct orientation in the wrong position or two vertices correct  A1 cao</p>



PAPER: 1MA0\_1H

Question		Working	Answer	Mark	Notes
24	(a)	$\overrightarrow{AB} = -\mathbf{a} + \mathbf{b}$ $\overrightarrow{ON} = \overrightarrow{OA} + \frac{2}{3}\overrightarrow{AB}$ $\overrightarrow{ON} = \mathbf{a} + \frac{2}{3}(-\mathbf{a} + \mathbf{b})$ $= \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$ OR $\overrightarrow{ON} = \overrightarrow{OB} + \frac{1}{3}\overrightarrow{BA}$ $\overrightarrow{ON} = \mathbf{b} + \frac{1}{3}(-\mathbf{b} + \mathbf{a})$ $= \frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$	$\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$	3	M1 for correct vector equation involving $\overrightarrow{ON}$ , eg. $\overrightarrow{ON} = \overrightarrow{OA} + \overrightarrow{AN}$ , may be written, partially or fully, in terms of $\mathbf{a}$ and $\mathbf{b}$ , e.g. ( $\overrightarrow{ON} =$ ) $\mathbf{a} + \frac{2}{3}\overrightarrow{AB}$ M1 for showing answer requires $\overrightarrow{AN} = \frac{2}{3}\overrightarrow{AB}$ or $\overrightarrow{BN} = \frac{1}{3}\overrightarrow{BA}$ A1 $\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}$ oe
	(b)	$\overrightarrow{OD} = \overrightarrow{OA} + \overrightarrow{AC} + \overrightarrow{CD}$ $= \mathbf{a} + \mathbf{b} + \mathbf{b}$ $= \mathbf{a} + 2\mathbf{b}$ $\overrightarrow{OD} = 3\left(\frac{1}{3}\mathbf{a} + \frac{2}{3}\mathbf{b}\right)$ $\overrightarrow{OD} = 3\overrightarrow{ON}$	Proof	3	M1 for a correct vector statement for $\overrightarrow{OD}$ or $\overrightarrow{ND}$ in terms of $\mathbf{a}$ and $\mathbf{b}$ , e.g. $\overrightarrow{OD} = \mathbf{a} + \mathbf{b} + \mathbf{b}$ oe or $\overrightarrow{ND} = \frac{2}{3}(-\mathbf{b} + \mathbf{a}) + \mathbf{b} + \mathbf{b}$ oe A1 for correct and fully simplified vectors for $\overrightarrow{ON}$ (may be seen in (a)) <b>and</b> for $\overrightarrow{OD} (= \mathbf{a} + 2\mathbf{b})$ or $\overrightarrow{ND} (= \frac{2}{3}\mathbf{a} + \frac{4}{3}\mathbf{b})$ C1 (dep on A1) for statement that $\overrightarrow{OD}$ or $\overrightarrow{ND}$ is a multiple of $\overrightarrow{ON}$ (+ common point)





Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

