Target Test Calculator 3

Mark Scheme

## NOTES ON MARKING PRINCIPLES

#### 1 Types of mark

M marks: method marks A marks: accuracy marks B marks: unconditional accuracy marks (independent of M marks)

#### 2 Abbreviations

cao - correct answer only ft - follow through isw - ignore subsequent working SC: special dep - dependent oe - or equivalent (and appropriate)

indep - independent

## 3 No working

If no working is shown then correct answers normally score full marks If no working is shown then incorrect (even though nearly correct) answers score no marks.

#### 4 With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks. Send the response to review, and discuss each of these situations with your Team Leader.

If there is no answer on the answer line then check the working for an obvious answer.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks. Discuss each of these situations with your Team Leader.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

## 5 Follow through marks

Follow through marks which involve a single stage calculation can be awarded without working since you can check the answer yourself, but if ambiguous do not award.

Follow through marks which involve more than one stage of calculation can only be awarded on sight of the relevant working, even if it appears obvious that there is only one way you could get the answer given.

## 6 Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: e.g. incorrect canceling of a fraction that would otherwise be correct

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect e.g. algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

# 7 Probability

Probability answers must be given a fractions, percentages or decimals. If a candidate gives a decimal equivalent to a probability, this should be written to at least 2 decimal places (unless tenths).

Incorrect notation should lose the accuracy marks, but be awarded any implied method marks.

If a probability answer is given on the answer line using both incorrect and correct notation, award the marks.

If a probability fraction is given then cancelled incorrectly, ignore the incorrectly cancelled answer.

## 8 Linear equations

Full marks can be gained if the solution alone is given on the answer line, or otherwise unambiguously indicated in working (without contradiction elsewhere). Where the correct solution only is shown substituted, but not identified as the solution, the accuracy mark is lost but any method marks can be awarded.

#### 9 Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

## 10 Range of answers

Unless otherwise stated, when an answer is given as a range (e.g 3.5 - 4.2) then this is inclusive of the end points (e.g 3.5, 4.2) and includes all numbers within the range (e.g 4, 4.1)

| 1380/4H  |   |        |      |   |  |
|----------|---|--------|------|---|--|
| Question | Working   | Answer | Mark | Notes   |  |
| 1        | 5 × 8 ÷ 2   | 20     | 2    | M1 for 5×8÷2 oe<br>A1 cao   |  |
| 2        | $ \begin{array}{rcl} 1 - 0.58 - 0.3 \\ = & 1 - 0.88 \end{array} $   | 0.12   | 2    | M1 for 1 – 0.58 – 0.3 oe<br>A1 for 0.12 oe  |  |
| 3        | $B = 20 \times 2 = 40$<br>$C = 3 \div 4 \times 20 = 15$<br>$D = 10 \div 100 \times 20 + 20 = 22$<br>20 + 40 + 15 + 22 | 97     | 4    | M1 for $20 \times 2$ or 40 seen<br>M1 for $3 \div 4 \times 20$ or 15 seen<br>M1 for $10 \div 100 \times 20 \div 20$ oe or 22 seen<br>or $1.1 \times 20$<br>A1 cao |  |
| 4 (a)    | 3 × 100   | 300    | 2    | M1 for 3 × 100 or 100 ÷ 6 × 18 oe<br>A1 cao   |  |
| (b)      | $2 \div \frac{1}{2} \times 6$   | 24     | 2    | M1 for $2 \div \frac{1}{2} \times 6$ oe<br>A1 cao   |  |

| 1380/4H  |                                   |                 |      |  |  |
|----------|-----------------------------------|-----------------|------|--|--|
| Question | Working                           | Answer          | Mark | Notes  |  |
| 5 (a)    |                                   |                 | 2    | B2 cao<br>(B1 for shape in the correct orientation<br>above the line $y = x$<br>or for shape elongated or shortened by one<br>square but with either top<br>or bottom in the correct position and in<br>the correct orientation)   |  |
| (b)      |                                   |                 | 3    | <ul> <li>B3 for correct enlargement in correct position</li> <li>(B2 for enlargement SF 3 in incorrect position or enlargement, centre <i>O</i>, but different scale factor)</li> <li>(B1 for 4 lines enlarged by SF 3 or enlargement, not from <i>O</i>, different scale factor)</li> </ul> |  |
| 6 (a)    |                                   | 6x + 5y         | 2    | B2<br>(B1 for either 6x or 5y seen)  |  |
| (b)      | 2x = 10 - 3 = 7<br>$x = 7 \div 2$ | 3.5             | 2    | M1 for $2x = 10 - 3$ or $2x = 7$<br>or $(10 - 3) \div 2$<br>A1 for 3.5 oe  |  |
| (c)(i)   |                                   | C <sup>11</sup> | 2    | B1 accept c <sup>5+6</sup>   |  |
| (ii)     |                                   | e <sup>8</sup>  |      | B1 accept e <sup>12-4</sup>  |  |

| 1380/4H |     |              |          |      |   |  |
|---------|-----|--------------|----------|------|---|--|
| Questic | on  | Working      | Answer   | Mark | Notes   |  |
| 7       |     | 8 ÷ 20 × 100 | 40       | 2    | M1 for $8 \div 20 \times 100$ or $\frac{8}{20} = \frac{8 \times 5}{20 \times 5}$ oe or<br>A1 cao  |  |
| 8 (     | (a) |              | 10 to 19 | 1    | B1 cao  |  |
|         | (b) |              | 20 to 29 | 1    | B1 for acceptable reason eg correct answer is 20<br>to 29<br>eg 30 <sup>th</sup> /31 <sup>st</sup> person not in this interval  |  |
|         | (c) |              |          | 2    | <ul> <li>B2 for complete polygon (ignore histograms and any lines below an age of 4.5 or above an age of 65, but award B1 if there is a line joining the first to the last point.)</li> <li>(B1 for one vertical or horizontal error OR incorrect but consistent error in placing the midpoints horizontally OR correct plotting but not joined. In this case ignore a line joining the first to the last.)</li> <li>Plotting tolerance: ± 1 square Points to be joined by lines (ruled or handdrawn, but not curves.)</li> </ul> |  |

| 1380/4H  |     |         |  |      |   |  |
|----------|-----|---------|--|------|---|--|
| Question |     | Working | Answer                                   | Mark | Notes   |  |
| 9        |     |         | 2.42927(0474)                            | 2    | B2 for 2.42927 or better<br>(B1 for 19.56 or 8.0518 seen or 2.43 or 2.429<br>or 2.4292 or 2.4293 or digits 242927<br>or seen) |  |
| 10       | (a) |         | -2, -1, 0, 1, 2                          | 2    | B2 for -2, -1, 0, 1, 2<br>(B1 for one extra or one missing)   |  |
|          | (b) | 2x < 30 | <i>x</i> < 15                            | 2    | M1 for $2x < 30$ or $\frac{x}{3} < 5$ or $x = 15$ or $x > 15$<br>A1 cao   |  |
| 11       |     |         | A and 3<br>B and 2<br>C and 4<br>D and 1 | 2    | B2 for all 4 correct<br>(B1 for 2 correct)  |  |

| 1380/4 | 1380/4H |   |                           |      |  |  |  |
|--------|---------|---|---------------------------|------|--|--|--|
| Ques   | stion   | Working   | Answer                    | Mark | Notes  |  |  |
| 12     | (a)     |   | (0, 3, 2)                 | 1    | B1 cao   |  |  |
|        | (b)     |   | (5, 3, 0)                 | 1    | B1 cao   |  |  |
| 13     | (a)     | 8.25 × 10 <sup>7</sup>                              | 8.25 × 10 <sup>7</sup>    | 1    | B1 cao   |  |  |
|        | (b)     | = 14. 56 × 10 <sup>-16</sup>                        | 1.456 × 10 <sup>-15</sup> | 2    | M1 for digits $1456 \times 10^{n}$ or $A \times 10^{-15}$ , $1 < A < 2$<br>A1 for $1.456 \times 10^{-15}$  |  |  |
| 14     | (a)     | 2) 56<br>2) 28<br>2) 14<br>7                        | 2 × 2 × 2 × 7             | 2    | M1 for a systematic method of at least 2<br>correct divisions<br>by a prime number oe factor tree, can be<br>implied by<br>digits 2, 2, 2, 7 on answer line<br>A1 for $2 \times 2 \times 2 \times 7$ or $2^3 \times 7$   |  |  |
|        | (b)     | $42 = 2 \times 3 \times 7$                          | 14                        | 2    | <ul> <li>M1 for attempt to find the common factor<br/>(by 2 lists or otherwise) or for answer of 7</li> <li>A1 cao</li> </ul>  |  |  |
| 15     |         | $AB = 8 \cos 37^{\circ} = 8$<br>(0.7986)<br>= 6.389 | 6.39                      | 3    | M1 for cos37 =<br>M1 for $AB = 8 \cos 37^{\circ}$ or 6.4 seen (dep on 1 <sup>st</sup><br>M1)<br>A1 for 6.38 - 6.39<br>OR<br>M1 for $\frac{AB}{Sin53} = \frac{8}{Sin90}$<br>M1 for $AB = \frac{8Sin53}{Sin90} AB$<br>or 6.4 seen (dep on 1 <sup>st</sup> M1)<br>A1 for for 6.38 - 6.39<br>SC M2A0 for 6.12 (radians) or 6.69 (grad) |  |  |

| 1380/4H |       |         |                           |      |   |
|---------|-------|---------|---------------------------|------|---|
| Que     | stion | Working | Answer                    | Mark | Notes   |
| 16      | (a)   |         | 61, 82, 94, 100           | 1    | B1 cao  |
|         | (b)   |         | Points plotted and joined | 2    | <ul> <li>B2 ft (dep on sensible table - condone 1 addition error) for 5 points plotted correctly, ± 1 square, at ends of interval and joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points</li> <li>(B1 ft for 4 points plotted correctly and joined or for 5 points plotted correctly)</li> <li>(SC B1 if 5 points plotted not at end but consistent within each interval and joined)</li> </ul> |

| 1380/4H  | 1380/4H   |                               |      |  |  |  |  |
|----------|---|-------------------------------|------|--|--|--|--|
| Question | Working   | Answer                        | Mark | Notes  |  |  |  |
| 17 (a)   |   | -15, (-8), -7, -6, 1,<br>(20) | 2    | B2 for all 4 correct<br>(B1 for 2 or 3 correct)  |  |  |  |
| (b)      |   |                               | 2    | <ul> <li>B2 for fully correct graph</li> <li>OR</li> <li>B1 ft for 6 'points' plotted correctly ± 1square</li> <li>B1 for smooth curve through all their 5 or 6</li> <li>plotted points provided B1 awarded in (a)</li> </ul>  |  |  |  |
| 18       | $\frac{7}{11} \times \frac{4}{10} + \frac{4}{11} \times \frac{7}{10}$ $= \frac{28}{110} + \frac{28}{110}$ | $\frac{28}{55}$               | 3    | M1 for $\frac{4}{10}$ and $\frac{7}{10}$ as second probabilities,<br>may be seen on a tree diagram, or for $\frac{7}{11} \times \frac{4}{10}$ or $\frac{4}{11} \times \frac{7}{10}$<br>M1 (dep) for $\times \frac{4}{10} + \frac{4}{11} \times \frac{7}{10}$<br>A1 for oe<br>SC B2 for an answer of oe |  |  |  |